

产品名称: MAP1A Rabbit Polyclonal Antibody
产品货号: APRab13622



产品概述 (Summary)

产品名称 (Production Name)	MAP1A Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MAP1A
别名 (Alternative Names)	MAP1L
基因 ID (Gene ID)	4130.0
蛋白 ID (SwissProt ID)	P78559. Synthesized peptide derived from human protein . at AA range: 1860-1940

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:300, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	308kDa

研究背景 (Background)

This gene encodes a protein that belongs to the microtubule-associated protein family. The proteins of this family are

产品名称: MAP1A Rabbit Polyclonal Antibody
产品货号: AP Rab13622




thought to be involved in microtubule assembly, which is an essential step in neurogenesis. The product of this gene is a precursor polypeptide that presumably undergoes proteolytic processing to generate the final MAP1A heavy chain and LC2 light chain. Expression of this gene is almost exclusively in the brain. Studies of the rat microtubule-associated protein 1A gene suggested a role in early events of spinal cord development. [provided by RefSeq, Jul 2008],domain:The basic region containing the repeats may be responsible for the binding of MAP1A to microtubules.,function:Structural protein involved in the filamentous cross-bridging between microtubules and other skeletal elements.,PTM:LC2 is generated from MAP1A by proteolytic processing.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MAP1 family.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins. Interacts with TIAM2. Interacts with guanylate kinase-like domain of DLG1, DLG2, DLG4.,tissue specificity:Brain.,

研究领域 (Research Area)

Signal Transduction; Cytoskeleton / ECM; Cytoskeleton; Microtubules; MT Associated Proteins; MAP; Neuroscience; Neurology process; Neurogenesis

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200 (4° overnight) .
2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only .