

产品名称: MAP1B Rabbit Polyclonal Antibody
产品货号: AP Rab13623



产品概述 (Summary)

产品名称 (Production Name)	MAP1B Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MAP1B
别名 (Alternative Names)	MAP1B
基因 ID (Gene ID)	4131.0
蛋白 ID (SwissProt ID)	P46821. Synthesized peptide derived from human protein . at AA range: 480-560

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:300, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	271kDa

研究背景 (Background)

This gene encodes a protein that belongs to the microtubule-associated protein family. The proteins of this family are

产品名称: MAP1B Rabbit Polyclonal Antibody
产品货号: AP Rab13623




thought to be involved in microtubule assembly, which is an essential step in neurogenesis. The product of this gene is a precursor polypeptide that presumably undergoes proteolytic processing to generate the final MAP1B heavy chain and LC1 light chain. Gene knockout studies of the mouse microtubule-associated protein 1B gene suggested an important role in development and function of the nervous system. [provided by RefSeq, Jul 2008],domain:Has a highly basic region with many copies of the sequence KKEE and KKEI/V, repeated but not at fixed intervals, which is responsible for the binding of MAP1B to microtubules.,function:The function of brain MAPS is essentially unknown. Phosphorylated MAP1B may play a role in the cytoskeletal changes that accompany neurite extension. Possibly MAP1B Binds to at least two tubulin subunits in the polymer, and this bridging of subunits might be involved in nucleating microtubule polymerization and in stabilizing microtubules.,PTM:LC1 is generated from MAP1B by proteolytic processing.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MAP1 family.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins. LC1 interacts with the amino-terminal region of MAP1B. Interacts with ANP32A and TIAM2 (By similarity). Interacts (via C-terminus) with GAN (via Kelch domains),.

研究领域 (Research Area)

Neurology process; Neurogenesis; Neuroscience; Cell Type Marker; Neuron marker; Axon marker; Neuroscience Markers; Neuronal; Cell Adhesion Proteins; Cytoskeletal Proteins; Microtubules; Signal Transduction; Cytoskeleton / ECM; Cytoskeleton; Microtubules; MT Associated Proteins; MAP; Tags & Cell Markers

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only .