

产品名称: MASP-2 Rabbit Polyclonal Antibody
产品货号: APRab13655



产品概述 (Summary)

产品名称 (Production Name)	MASP-2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MASP2
别名 (Alternative Names)	MASP2; Mannan-binding lectin serine protease 2; MBL-associated serine protease 2; Mannose-binding protein-associated serine protease 2; MASP-2
基因 ID (Gene ID)	10747.0
蛋白 ID (SwissProt ID)	O00187.The antiserum was produced against synthesized peptide derived from human MASP2. AA range:227-276

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	75kDa

产品名称: MASP-2 Rabbit Polyclonal Antibody
产品货号: APRab13655




研究背景 (Background)

mannan binding lectin serine peptidase 2(MASP2) Homo sapiens This gene encodes a member of the peptidase S1 family of serine proteases. The encoded preproprotein is proteolytically processed to generate A and B chains that heterodimerize to form the mature protease. This protease cleaves complement components C2 and C4 in order to generate C3 convertase in the lectin pathway of the complement system. The encoded protease also plays a role in the coagulation cascade through cleavage of prothrombin to form thrombin. Myocardial infarction and acute stroke patients exhibit reduced serum concentrations of the encoded protein. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016],catalytic activity:Selective cleavage after Arg-223 in complement component C2 (-Ser-Leu-Gly-Arg-|-Lys-Ile-Gln-Ile) and after Arg-76 in complement component C4 (-Gly-Leu-Gln-Arg-|-Ala-Leu-Glu-Ile),disease:Genetic variation in MASP2 is the cause of MASP2 deficiency [MIM:605102]. It is associated with susceptibility to infections and with the development of immunologic disease.,function:Serum protease that plays an important role in the activation of the complement system via mannose-binding lectin. After activation by auto-catalytic cleavage it cleaves C2 and C4, leading to their activation and to the formation of C3 convertase.,miscellaneous:Dimerization and MBL2 binding requires calcium ions.,online information:MASP2 mutation db,PTM:Activated by cleavage after Arg-444. The uncleaved zymogen is inactive towards synthetic substrates, but has sufficient activity to effect autocatalytic cleavage.,PTM:The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 EGF-like domain.,similarity:Contains 1 peptidase S1 domain.,similarity:Contains 2 CUB domains.,similarity:Contains 2 Sushi (CCP/SCR) domains.,subunit:Homodimer; disulfide-linked. Binds MBL2. Isoform 2 binds to MASP1. Binds SERPING1.,tissue specificity:Plasma.,

研究领域 (Research Area)


Complement and coagulation cascades;

图片 (Image Data)




Western blot analysis of MASP2 Antibody. The lane on the right is blocked with the MASP2 peptide.

产品名称: MASP-2 Rabbit Polyclonal Antibody
产品货号: APRab13655



Western blot analysis of the lysates from HT-29 cells using MASP2 antibody.



Western Blot analysis of K562 cells using MASP-2 Polyclonal Antibody

注意事项 (Note)

For research use only .