

产品名称: MEF-2C Rabbit Polyclonal Antibody
产品货号: APRab13786



产品概述 (Summary)

产品名称 (Production Name)	MEF-2C Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MEF2C
别名 (Alternative Names)	MEF2C; Myocyte-specific enhancer factor 2C
基因 ID (Gene ID)	4208.0
蛋白 ID (SwissProt ID)	Q06413.The antiserum was produced against synthesized peptide derived from human MEF2C. AA range:362-411

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	51kDa

研究背景 (Background)

产品名称: MEF-2C Rabbit Polyclonal Antibody
产品货号: AP Rab13786




This locus encodes a member of the MADS box transcription enhancer factor 2 (MEF2) family of proteins, which play a role in myogenesis. The encoded protein, MEF2 polypeptide C, has both trans-activating and DNA binding activities. This protein may play a role in maintaining the differentiated state of muscle cells. Mutations and deletions at this locus have been associated with severe mental retardation, stereotypic movements, epilepsy, and cerebral malformation. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jul 2010], alternative products: Additional isoforms seem to exist, developmental stage: Expression is highest during the early stages of postnatal development, at later stages levels greatly decrease., domain: The beta domain, missing in a number of isoforms, is required for enhancement of transcriptional activity., function: Transcription activator which binds specifically to the MEF2 element present in the regulatory regions of many muscle-specific genes. Controls cardiac morphogenesis and myogenesis, and is also involved in vascular development. May also be involved in neurogenesis and in the development of cortical architecture (By similarity). Isoform 3 and isoform 4, which lack the repressor domain, are more active than isoform 1 and isoform 2., PTM: Acetylated by p300 on several sites in differentiating myocytes. Acetylation on Lys-4 increases DNA binding and transactivation., PTM: Phosphorylation on Ser-59 enhances DNA binding activity (By similarity). Phosphorylation on Ser-396 is required for Lys-391 sumoylation and inhibits transcriptional activity., PTM: Proteolytically cleaved in cerebellar granule neurons, probably by caspase 7, following neurotoxicity. Preferentially cleaves the CDK5-mediated hyperphosphorylated form which leads to neuron apoptosis and transcriptional inactivation., PTM: Sumoylated on Lys-391 by SUMO2 but not by SUMO1 represses transcriptional activity., similarity: Belongs to the MEF2 family., similarity: Contains 1 MADS-box domain., similarity: Contains 1 Mef2-type DNA-binding domain., subunit: Forms a complex with class II HDACs in undifferentiating cells. On myogenic differentiation, HDACs are released into the cytoplasm allowing MEF2s to interact with other proteins for activation. Interacts with EP300 in differentiating cells; the interaction acetylates MEF2C leading to increased DNA binding and activation. Interacts with HDAC7 and CARM1 (By similarity). Interacts with HDAC4, HDAC7 AND HDAC9; the interaction WITH HDACs represses transcriptional activity., tissue specificity: Expressed in brain and skeletal muscle.,

研究领域 (Research Area)


AMPK; Protein_Acetylation; MAPK_ERK_Growth; MAPK_G_Protein

图片 (Image Data)

产品名称: MEF-2C Rabbit Polyclonal Antibody
产品货号: APRab13786



Western blot analysis of lysates from NIH/3T3 cells, treated with starved 24h, using MEF2C Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of 293T cells using MEF-2C Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by MinuteTM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA) .

注意事项 (Note)

For research use only .