

产品名称: mGluR-4 Rabbit Polyclonal Antibody
产品货号: AP Rab13861



产品概述 (Summary)

产品名称 (Production Name)	mGluR-4 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	GRM4
别名 (Alternative Names)	GRM4; GPRC1D; MGLUR4; Metabotropic glutamate receptor 4; mGluR4
基因 ID (Gene ID)	2914.0
蛋白 ID (SwissProt ID)	Q14833.The antiserum was produced against synthesized peptide derived from human mGluR4. AA range:851-900

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	100kDa

研究背景 (Background)

产品名称: mGluR-4 Rabbit Polyclonal Antibody
产品货号: AP Rab13861




glutamate metabotropic receptor 4 (GRM4) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this function: Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity., similarity: Belongs to the G-protein coupled receptor 3 family., subunit: Interacts with PICK1., tissue specificity: Strongly expressed in the cerebellum. Expressed at low levels in hippocampus, hypothalamus and thalamus. No expression detected in liver.,


研究领域 (Research Area)

Neuroactive ligand-receptor interaction; Taste transduction;

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using mGluR4 Antibody. The picture on the right is blocked with the synthesized peptide.




产品名称: mGluR-4 Rabbit Polyclonal Antibody
产品货号: AP Rab13861



Western blot analysis of lysates from 293 cells, treated with Forskolin 40nM 30', using mGluR4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using mGluR-4 Polyclonal Antibody diluted at 1: 500



Western Blot analysis of 293 cells using mGluR-4 Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .