

产品名称: N2DL2 Rabbit Polyclonal Antibody
产品货号: APRab14366



产品概述 (Summary)

产品名称 (Production Name)	N2DL2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ULBP2 N2DL2 RAET1H UNQ463/PRO791
别名 (Alternative Names)	
基因 ID (Gene ID)	80328.0
蛋白 ID (SwissProt ID)	Q9BZM5.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	27kDa

研究背景 (Background)

This gene encodes a major histocompatibility complex (MHC) class I-related molecule that binds to the NKG2D receptor on natural killer (NK) cells to trigger release of multiple cytokines and chemokines that in turn contribute to the recruitment


产品名称: N2DL2 Rabbit Polyclonal Antibody
产品货号: AP Rab14366

and activation of NK cells. The encoded protein undergoes further processing to generate the mature protein that is either anchored to membrane via a glycosylphosphatidylinositol moiety, or secreted. Many malignant cells secrete the encoded protein to evade immunosurveillance by NK cells. This gene is located in a cluster of multiple MHC class I-related genes on chromosome 6. [provided by RefSeq, Jul 2015],function:Ligand for the NKG2D receptor, together with at least ULBP1 and ULBP3. ULBPs activate multiple signaling pathways in primary NK cells, resulting in the production of cytokines and chemokines. Binding of ULBPs ligands to NKG2D induces calcium mobilization and activation of the JAK2, STAT5, ERK and PI3K kinase/Akt signal transduction pathway. In CMV infected cells, interacts with soluble CMV glycoprotein UL16. The interaction with UL16 blocked the interaction with the NKG2D receptor, providing a mechanism by which CMV infected cells might escape the immune system. UL16 also causes ULBP2 to be retained in the ER and cis-Golgi apparatus so that it does not reach the cell surface.,miscellaneous:The ULBPs are unusual members of the extended MHC class I superfamily, because they do not contain the alpha 3 domain and they lack a transmembrane domain. They are unlikely to present peptides.,similarity:Belongs to the MHC class I family.,subunit:Interacts with the CMV glycoprotein UL16. Does not bind to beta2-microglobulin.,tissue specificity:Expressed in various types of cancer cell lines and in the fetus, but not in normal tissues.,

研究领域 (Research Area)

Natural killer cell mediated cytotoxicity;

图片 (Image Data)



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night

注意事项 (Note)

For research use only .