

产品名称: Nanog P8 Rabbit Polyclonal Antibody
产品货号: AP Rab14403



产品概述 (Summary)

产品名称 (Production Name)	Nanog P8 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NANOGP8
别名 (Alternative Names)	NANOGP8; Putative homeobox protein NANOGP8
基因 ID (Gene ID)	388112.0
蛋白 ID (SwissProt ID)	Q6NSW7.The antiserum was produced against synthesized peptide derived from human NANOGP8. AA range:51-100

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	35kDa

研究背景 (Background)

产品名称: Nanog P8 Rabbit Polyclonal Antibody
产品货号: AP Rab14403




This locus is a processed pseudogene of the transcription factor NANOG. NANOG plays a central role in regulating self-renewal in pluripotent stem cells and tumor cells. This pseudogene contains an intact open reading frame that could potentially encode a protein similar to NANOG. Although there is no evidence of transcription from this pseudogene, RT-PCR studies suggest that NANOGP8 may be expressed in some cancer cell lines. In vitro studies using a recombinant NANOGP8 protein have shown that the protein localizes to the nucleus and can promote cell proliferation, similar to NANOG. [provided by RefSeq, Sep 2009],developmental stage:Expressed in embryonic stem (ES) and carcinoma (EC) cells. Expressed in inner cell mass (ICM) of the blastocyst and gonocytes between 14 and 19 weeks of gestation (at protein level). Not expressed in oocytes, unfertilized oocytes, 2-16 cell embryos and early morula (at protein level). Expressed in embryonic stem cells (ES). Expression decreases with ES differentiation.,function:May act as a transcription regulator (By similarity). When overexpressed, promotes cells to enter into S phase and proliferation.,function:Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes (By similarity). Acts as a transcriptional activator or repressor (By similarity). Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3' (By similarity). When overexpressed, promotes cells to enter into S phase and proliferation.,miscellaneous:Almost identical to NANOG. There are only one change in the inferred amino acid sequence from 'Gln-253' in NANOG to His-253 in NANOGP8.,miscellaneous:Exists an other tandem duplicated non-processed pseudogene (NANOGP1) and 10 other NANOG-related nucleotide sequences located on different chromosomes, all of which are processed pseudogenes lacking introns (NANOGP2 to NANOGP11); except NANOGP8 which is a retrogene.,online information:Nanog entry,similarity:Belongs to the Nanog homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,subunit:Interacts with SMAD1 and SALL4.,tissue specificity:Expressed in osteosarcoma cancer cell line (at protein level) (Probable). Expressed in tumor uterine cervix, breast and urinary bladder tissues, and also osteosarcoma, hepatoma, and breast adenocarcinoma cancer cell lines.,tissue specificity:Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.,


研究领域 (Research Area)

图片 (Image Data)

产品名称: Nanog P8 Rabbit Polyclonal Antibody
产品货号: AP Rab14403



Western blot analysis of lysates from HUVEC, HT-29, HepG2, and Jurkat cells, using NANOGP8 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Nanog P8 Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only .