

**产品名称:** NCoA-3 Rabbit Polyclonal Antibody

**产品货号:** APRab14454

### 产品概述 (Summary)

|                   |                                                                                                   |
|-------------------|---------------------------------------------------------------------------------------------------|
| 描述(Description)   | Rabbit polyclonal Antibody                                                                        |
| 宿主(Host)          | Rabbit                                                                                            |
| 应用(Application)   | WB,IHC,ICC/IF,ELISA                                                                               |
| 种属反应(Reactivity)  | Human,Mouse,Rat                                                                                   |
| 偶联物(Conjugation)  | Unconjugated                                                                                      |
| 修饰(Modification)  | Unmodified                                                                                        |
| 同种型(Isotype)      | IgG                                                                                               |
| 克隆(Clonality)     | Polyclonal                                                                                        |
| 剂型(Form)          | Liquid                                                                                            |
| 保存条件(Storage)     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液(Buffer)      | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化 (Purification) | Affinity purification                                                                             |

### 产品应用(Application)

|                       |                                                                         |
|-----------------------|-------------------------------------------------------------------------|
| 稀释比(Dilution Ratio)   | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000 |
| 分子量(Molecular Weight) | 160kDa                                                                  |

### 抗原信息(Antigen Information)

|                       |                                                                                                                                                                                                                                                                                                                                                                 |
|-----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因名(Gene Name)        | NCOA3                                                                                                                                                                                                                                                                                                                                                           |
| 别名(Alternative Names) | NCOA3; AIB1; BHLHE42; RAC3; TRAM1; Nuclear receptor coactivator 3; NCoA-3; ACTR; Amplified in breast cancer 1 protein; AIB-1; CBP-interacting protein; pCIP; Class E basic helix-loop-helix protein 42; bHLHe42; Receptor-associated coactivator 3; RAC-3; Steroid receptor coactivator protein 3; SRC-3; Thyroid hormone receptor activator molecule 1; TRAM-1 |
| 基因 ID(Gene ID)        | 8202.0                                                                                                                                                                                                                                                                                                                                                          |
| SwissProt ID          | Q9Y6Q9                                                                                                                                                                                                                                                                                                                                                          |
| 免疫原(Immunogen)        | Synthesized peptide derived from the Internal region of human NCoA-3.                                                                                                                                                                                                                                                                                           |

### 研究背景 (Background)

The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits p300/CBP-associated factor and CREB binding protein as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Several transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein. [provided by RefSeq, Mar 2010], alternative products: Additional isoforms seem to exist, catalytic activity: Acetyl-CoA + histone = CoA + acetylhistone, domain: Contains three Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. Motifs 1 and 2 are essential for the association with nuclear receptors, and constitute the RID domain (Receptor-interacting domain), enzyme regulation: Coactivator activity on nuclear receptors and NF-kappa-B pathways is enhanced by various hormones, and the TNF cytokine, respectively. TNF stimulation probably enhances phosphorylation, which in turn activates coactivator function. In contrast, acetylation by CREBBP apparently suppresses coactivation of target genes by disrupting its association with nuclear receptors, function: Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit, miscellaneous: NCOA3 is frequently amplified or overexpressed in breast and ovarian cancers, polymorphism: The length of the poly-Gln region is polymorphic in the normal population, PTM: Acetylated by CREBBP. Acetylation occurs in the RID domain, and disrupts the interaction with nuclear receptors and regulates its function, PTM: Methylated by CARM1, PTM: Phosphorylated by IKK complex. Regulated its function, similarity: Belongs to the SRC/p160 nuclear receptor coactivator family, similarity: Contains 1 basic helix-loop-helix (bHLH) domain, similarity: Contains 1 PAS (PER-ARNT-SIM) domain, subcellular location: Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus, subunit: Interacts with CARM1 (By similarity). Present in a complex containing NCOA2, IKKA, IKKB, IKBKG and the histone acetyltransferase protein CREBBP. Interacts with CASP8AP2, NR3C1 and PCAF. Interacts with ATAD2 and this interaction is enhanced by estradiol, tissue specificity: Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.


### 研究领域 (Research Area)

Pathways and Processes; Metabolic signaling pathways; Lipid and lipoprotein metabolism; Lipid metabolism; Metabolism; Types of disease; Obesity; Metabolism of lipids and lipoproteins; Signal transduction; Nuclear signaling; Nuclear hormone receptors; Coactivators and repressors; Cancer; Cancer Metabolism; Nuclear Signaling Pathways; Nuclear Receptors; Co-activators/co-repressors; Epigenetics and Nuclear Signaling; Chromatin Modifying Enzymes; Acetylation; HAT; Nuclear Signaling Pathways; Nuclear Receptors; Co-activators/co-repressors; Signaling Pathway; Nuclear Signaling


### 注意事项 (Note)

For research use only.

## 图片 (Image Data)



Western Blot analysis of HeLa, HepG2 cells using NCoA-3 Polyclonal Antibody..  
Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was  
diluted at 1:100

## EnkiLife 优势产品推荐

| 产品类别                        | 产品货号     | 产品名称                                               |
|-----------------------------|----------|----------------------------------------------------|
| WB 解决方案                     | RA10020  | 2h 极速 WB 即用型全流程试剂盒                                 |
|                             | RA10021  | 4h 快速 WB 即用型全流程试剂盒                                 |
|                             | RA10042  | 5h 畅享版 WB 全流程试剂盒                                   |
|                             | RA10037  | 校准级彩色预染蛋白 Marker (8-180kDa)                        |
|                             | RA10038  | 校准级彩色预染蛋白 Marker (10-250kDa)                       |
|                             | RA10039  | 校准级高分子彩色预染蛋白 Marker (25-400kDa)                    |
| TSA 多重荧光染色试剂盒               | RA10008  | TSA 双标三色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10009  | TSA 三标四色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10010  | TSA 四标五色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10011  | TSA 五标六色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10012  | TSA 六标七色多重荧光染色试剂盒 (mIHC)                           |
| IHC 检测试剂盒                   | RA10006  | HRP Anti-Mouse/Rabbit IHC Detection System         |
|                             | RA10007  | Polymer-HRP Anti-Mouse/Rabbit IHC Detection System |
| 抗体标记试剂盒                     | RE80004p | 辣根过氧化物酶(HRP)抗体标记试剂盒                                |
|                             | RE80002q | Sulfo-NHS-生物素标记试剂盒                                 |
|                             | RE80007p | Cy3 荧光素标记试剂盒                                       |
|                             | RE80011p | Fluor488 荧光素标记试剂盒                                  |
|                             | RE80017p | Fluor750 荧光素标记试剂盒                                  |
|                             | RE80005p | 藻红蛋白(R-PE) 抗体快速标记试剂盒                               |
|                             | RE80040  | PE-Cy7 串联染料抗体快速标记试剂盒                               |
| 稳转细胞系构建服务<br>(免费赠送全膜 WB 验证) | TS-0001  | 过表达稳转细胞系构建                                         |
|                             | TS-0002  | 敲低稳转细胞系构建                                          |
|                             | TS-0003  | 敲除细胞系构建                                            |

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>