

产品名称: N-CoR Rabbit Polyclonal Antibody

产品货号: APRab14457

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Mouse
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
分子量(Molecular Weight)	270kDa

抗原信息(Antigen Information)

基因名(Gene Name)	NCOR1
别名(Alternative Names)	NCOR1; KIAA1047; Nuclear receptor corepressor 1; N-CoR; N-CoR1
基因 ID(Gene ID)	9611.0
SwissProt ID	O75376
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human NCoR1. AA range:51-100

研究背景 (Background)

This gene encodes a protein that mediates ligand-independent transcription repression of thyroid-hormone and retinoic-acid receptors by promoting chromatin condensation and preventing access of the transcription machinery. It is part of a complex

which also includes histone deacetylases and transcriptional regulators similar to the yeast protein Sin3p. This gene is located between the Charcot-Marie-Tooth and Smith-Magenis syndrome critical regions on chromosome 17. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 17 and 20.[provided by RefSeq, Jun 2010],domain:The C-terminal region contains two separate nuclear receptor-interacting domains (ID1 and ID2), each of which contains a conserved sequence referred to as the CORNR box. This motif is necessary and sufficient for binding to unligated nuclear hormone receptors, while sequences flanking the CORNR box determine the precise nuclear hormone receptor specificity.,domain:The N-terminal region contains three independent domains that are capable of mediating transcriptional repression (RD1, RD2 and RD3),function:Mediates transcriptional repression by certain nuclear receptors. Part of a complex which promotes histone deacetylation and the formation of repressive chromatin structures which may impede the access of basal transcription factors.,PTM:Ubiquitinated; mediated by SIAH2 and leading to its subsequent proteasomal degradation.,similarity:Belongs to the N-CoR nuclear receptor corepressors family.,similarity:Contains 2 SANT domains.,subunit:Interacts with SIAH2, HDAC7, SAP30, SIN3A and SIN3B (By similarity). Forms a large corepressor complex that contains SIN3A/B and histone deacetylases HDAC1 and HDAC2. This complex associates with the thyroid (TR) and the retinoid acid receptors (RAR) in the absence of ligand. Interacts with DACH1. Component of the N-CoR repressor complex, at least composed of NCOR1, NCOR2, HDAC3, TBL1X, TBL1XR1, CORO2A and GPS2. Interacts with TRIM28 and MJD2A/JHDM3A. Interacts with ZBTB33/KAISO, and this interaction serves to recruit the N-CoR complex to promoter regions containing methylated CpG dinucleotides. Interacts with the catalytic domain of HDAC9. Interacts with CBFA2T3 and HEXIM1. Interacts with C1D.,


研究领域 (Research Area)

Epigenetics and Nuclear Signaling; Chromatin Modifying Enzymes; Acetylation; Signal Transduction; Signaling Pathway; Nuclear Signaling; Nuclear Hormone Receptors; Co-activators/co-repressors; Nuclear Signaling Pathways; Nuclear Receptors; Transcription; Co-factors; Acetylation; HDACs; Class II / Hda1 Class; Cancer; Cancer Metabolism; Metabolic signaling pathway; Metabolism of lipids and lipoproteins; Metabolism; Pathways and Processes; Lipid and lipoprotein metabolism; Lipid metabolism; Cofactors, Vitamins / minerals


注意事项 (Note)

For research use only.

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using NCoR1 Antibody. The picture on the right is blocked with the synthesized peptide.



EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>