

产品名称: Nek9 Rabbit Polyclonal Antibody
产品货号: AP Rab14559



产品概述 (Summary)

产品名称 (Production Name)	Nek9 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NEK9 NEK9; KIAA1995; NEK8; NERCC; Serine/threonine-protein kinase Nek9; Nercc1
别名 (Alternative Names)	kinase; Never in mitosis A-related kinase 9; NimA-related protein kinase 9; NimA-related kinase 8; Nek8
基因 ID (Gene ID)	91754.0
蛋白 ID (SwissProt ID)	Q8TD19. The antiserum was produced against synthesized peptide derived from human NEK9. AA range: 176-225

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, ICC/IF 1:200-1:1000, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	107kDa

产品名称: Nek9 Rabbit Polyclonal Antibody
产品货号: AP Rab14559




研究背景 (Background)

This gene encodes a member of the NimA (never in mitosis A) family of serine/threonine protein kinases. The encoded protein is activated in mitosis and, in turn, activates other family members during mitosis. This protein also mediates cellular processes that are essential for interphase progression. [provided by RefSeq, Jul 2016], catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., developmental stage: Expression varied mildly across the cell cycle, with highest expression observed in G1 and stationary-phase cells., domain: Dimerizes through its coiled-coil domain., enzyme regulation: Activated during mitosis by intramolecular autophosphorylation. Activity and autophosphorylation is activated by manganese >> magnesium ions. Sensitive to increasing concentration of detergents. It is not cell-cycle regulated but activity is higher in G0-arrested cells., function: Pleiotropic regulator of mitotic progression, participating in the control of spindle dynamics and chromosome separation. Phosphorylates different histones, myelin basic protein, beta-casein, and BICD2. Phosphorylates histone H3 on serine and threonine residues and beta-casein on serine residues. Important for G1/S transition and S phase progression., PTM: Autophosphorylated on serine and threonine residues. When complexed with FACT, exhibits markedly elevated phosphorylation on Thr-210. During mitosis, not phosphorylated on Thr-210. Phosphorylated by CDC2 in vitro., similarity: Belongs to the protein kinase superfamily. NEK Ser/Thr protein kinase family. NIMA subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 6 RCC1 repeats., subunit: Homodimer. Binds to Ran GTPase. Has a greater affinity for Ran-GDP over Ran-GTP. Interacts with NEK6, NEK7 and BICD2. Interacts with SSRP1 and SUPT16H, the 2 subunits of the FACT complex., tissue specificity: Most abundant in heart, liver, kidney and testis. Also expressed in smooth muscle cells and fibroblasts.,

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



产品名称: Nek9 Rabbit Polyclonal Antibody
产品货号: APRab14559



注意事项 (Note)

For research use only .