

**产品名称:** NOS2 Rabbit Polyclonal Antibody

**产品货号:** APRab14803

### 产品概述 (Summary)

|                   |                                                                                                   |
|-------------------|---------------------------------------------------------------------------------------------------|
| 描述(Description)   | Rabbit polyclonal Antibody                                                                        |
| 宿主(Host)          | Rabbit                                                                                            |
| 应用(Application)   | WB,IHC,ICC/IF,ELISA                                                                               |
| 种属反应(Reactivity)  | Human,Mouse,Rat                                                                                   |
| 偶联物(Conjugation)  | Unconjugated                                                                                      |
| 修饰(Modification)  | Unmodified                                                                                        |
| 同种型(Isotype)      | IgG                                                                                               |
| 克隆(Clonality)     | Polyclonal                                                                                        |
| 剂型(Form)          | Liquid                                                                                            |
| 保存条件(Storage)     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液(Buffer)      | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化 (Purification) | Affinity purification                                                                             |

### 产品应用(Application)

|                       |                                                                        |
|-----------------------|------------------------------------------------------------------------|
| 稀释比(Dilution Ratio)   | WB 1:500-1:2000,IHC 1:50-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000 |
| 分子量(Molecular Weight) | 131kDa                                                                 |

### 抗原信息(Antigen Information)

|                       |                                                                                                                                                                       |
|-----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因名(Gene Name)        | NOS2                                                                                                                                                                  |
| 别名(Alternative Names) | NOS2; NOS2A; Nitric oxide synthase; inducible; Hepatocyte NOS; HEP-NOS; Inducible NO synthase; Inducible NOS; iNOS; NOS type II; Peptidyl-cysteine S-nitrosylase NOS2 |
| 基因 ID(Gene ID)        | 4843.0                                                                                                                                                                |
| SwissProt ID          | P35228                                                                                                                                                                |
| 免疫原(Immunogen)        | The antiserum was produced against synthesized peptide derived from human iNOS. AA range:117-166                                                                      |

### 研究背景 (Background)

Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission and

antimicrobial and antitumoral activities. This gene encodes a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. Three related pseudogenes are located within the Smith-Magenis syndrome region on chromosome 17. [provided by RefSeq, Jul 2008],catalytic activity:L-arginine + n NADPH + n H(+) + m O(2) = citrulline + nitric oxide + n NADP(+),cofactor: Binds 1 FAD,cofactor: Binds 1 FMN,cofactor: Heme group,cofactor: Tetrahydrobiopterin (BH4). May stabilize the dimeric form of the enzyme,enzyme regulation: Regulated by calcium/calmodulin. Aspirin inhibits expression and function of this enzyme and effects may be exerted at the level of translational/post-translational modification and directly on the catalytic activity,function: Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal actions,induction: By endotoxins and cytokines,online information: Nitric oxide synthase entry,similarity: Belongs to the NOS family,similarity: Contains 1 FAD-binding FR-type domain,similarity: Contains 1 flavodoxin-like domain,subunit: Homodimer. Binds SLC9A3R1,tissue specificity: Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets,


### 研究领域 (Research Area)

Arginine and proline metabolism;Calcium;Pathways in cancer;Small cell lung cancer;


### 注意事项 (Note)

For research use only.


### 图片 (Image Data)




Western blot analysis of lysates from NIH/3T3 cells, using iNOS Antibody. The lane on the right is blocked with the synthesized peptide.




Immunofluorescence analysis of rat-kidney tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of rat-kidney tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of mouse-lung tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of mouse-lung tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of mouse-kidney tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of mouse-kidney tissue. 1, NOS2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human-uterus-cancer tissue. 1, NOS2 Polyclonal Antibody was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, Secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.



Western Blot analysis of various cells using NOS2 Polyclonal Antibody diluted at 1:500

## EnkiLife 优势产品推荐

| 产品类别                        | 产品货号     | 产品名称                                               |
|-----------------------------|----------|----------------------------------------------------|
| WB 解决方案                     | RA10020  | 2h 极速 WB 即用型全流程试剂盒                                 |
|                             | RA10021  | 4h 快速 WB 即用型全流程试剂盒                                 |
|                             | RA10042  | 5h 畅享版 WB 全流程试剂盒                                   |
|                             | RA10037  | 校准级彩色预染蛋白 Marker (8-180kDa)                        |
|                             | RA10038  | 校准级彩色预染蛋白 Marker (10-250kDa)                       |
|                             | RA10039  | 校准级高分子彩色预染蛋白 Marker (25-400kDa)                    |
| TSA 多重荧光染色试剂盒               | RA10008  | TSA 双标三色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10009  | TSA 三标四色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10010  | TSA 四标五色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10011  | TSA 五标六色多重荧光染色试剂盒 (mIHC)                           |
|                             | RA10012  | TSA 六标七色多重荧光染色试剂盒 (mIHC)                           |
| IHC 检测试剂盒                   | RA10006  | HRP Anti-Mouse/Rabbit IHC Detection System         |
|                             | RA10007  | Polymer-HRP Anti-Mouse/Rabbit IHC Detection System |
| 抗体标记试剂盒                     | RE80004p | 辣根过氧化物酶(HRP)抗体标记试剂盒                                |
|                             | RE80002q | Sulfo-NHS-生物素标记试剂盒                                 |
|                             | RE80007p | Cy3 荧光素标记试剂盒                                       |
|                             | RE80011p | Fluor488 荧光素标记试剂盒                                  |
|                             | RE80017p | Fluor750 荧光素标记试剂盒                                  |
|                             | RE80005p | 藻红蛋白(R-PE) 抗体快速标记试剂盒                               |
|                             | RE80040  | PE-Cy7 串联染料抗体快速标记试剂盒                               |
| 稳转细胞系构建服务<br>(免费赠送全膜 WB 验证) | TS-0001  | 过表达稳转细胞系构建                                         |
|                             | TS-0002  | 敲低稳转细胞系构建                                          |
|                             | TS-0003  | 敲除细胞系构建                                            |

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>