

产品名称: Nrf2 Rabbit Polyclonal Antibody
产品货号: AP Rab14893



产品概述 (Summary)

产品名称 (Production Name)	Nrf2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NFE2L2
别名 (Alternative Names)	NFE2L2; NRF2; Nuclear factor erythroid 2-related factor 2; NF-E2-related factor 2; NFE2-related factor 2; HEBP1; Nuclear factor; erythroid derived 2, like 2
基因 ID (Gene ID)	4780.0
蛋白 ID (SwissProt ID)	Q16236.The antiserum was produced against synthesized peptide derived from human Nrf2. AA range:556-605

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	75-100kDa

产品名称: Nrf2 Rabbit Polyclonal Antibody
产品货号: AP Rab14893




研究背景 (Background)

This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflammation which includes the production of free radicals. Multiple transcript variants encoding different isoforms have been characterized for this gene. [provided by RefSeq, Sep 2015], domain: Acidic activation domain in the N-terminus, and DNA binding domain in the C-terminus., function: Transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. Important for the coordinated up-regulation of genes in response to oxidative stress. May be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region., PTM: Phosphorylation of Ser-40 by PKC in response to oxidative stress dissociates NFE2L2 from its cytoplasmic inhibitor KEAP1, promoting its translocation into the nucleus., similarity: Belongs to the bZIP family., similarity: Belongs to the bZIP family. CNC subfamily., similarity: Contains 1 bZIP domain., subcellular location: Cytosolic under unstressed conditions, translocates into the nucleus upon induction by electrophilic agents., subunit: Heterodimer. May bind DNA with an unknown protein. Interacts with KEAP1. Interacts via its leucine-zipper domain with the coiled-coil domain of PMF1., tissue specificity: Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.,

研究领域 (Research Area)


Epigenetics and Nuclear Signaling

图片 (Image Data)




Immunofluorescence analysis of HUVEC cells, using Nrf2 Antibody. The picture on the right is blocked with the synthesized peptide.


产品名称: Nrf2 Rabbit Polyclonal Antibody
产品货号: APRab14893



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using Nrf2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC cells, using Nrf2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Nrf2 Polyclonal Antibody diluted at 1: 1000

产品名称: Nrf2 Rabbit Polyclonal Antibody
产品货号: APRab14893



Western Blot analysis of HuvEc cells using Nrf2 Polyclonal Antibody diluted at 1 : 1000

注意事项 (Note)

For research use only .