

产品名称: PAF-R Rabbit Polyclonal Antibody
产品货号: APRab15693



产品概述 (Summary)

产品名称 (Production Name)	PAF-R Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PTAFR
别名 (Alternative Names)	PTAFR; PAFR; Platelet-activating factor receptor; PAF-R; PAFr
基因 ID (Gene ID)	5724.0
蛋白 ID (SwissProt ID)	P25105.The antiserum was produced against synthesized peptide derived from human PTAFR. AA range:194-243

产品应用 (Application)

稀释比 (Dilution Ratio)	ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)

产品名称: PAF-R Rabbit Polyclonal Antibody
产品货号: AP Rab15693




This gene encodes a seven-transmembrane G-protein-coupled receptor for platelet-activating factor (PAF) that localizes to lipid rafts and/or caveolae in the cell membrane. PAF (1-0-alkyl-2-acetyl-sn-glycero-3-phosphorylcholine) is a phospholipid that plays a significant role in oncogenic transformation, tumor growth, angiogenesis, metastasis, and pro-inflammatory processes. Binding of PAF to the PAF-receptor (PAFR) stimulates numerous signal transduction pathways including phospholipase C, D, A2, mitogen-activated protein kinases (MAPKs), and the phosphatidylinositol-calcium second messenger system. Following PAFR activation, cells become rapidly desensitized and this refractory state is dependent on PAFR phosphorylation, internalization, and down-regulation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2011],function:Receptor for platelet activating factor, a chemotactic phospholipid mediator that possesses potent inflammatory, smooth-muscle contractile and hypotensive activity. Seems to mediate its action via a G protein that activates a phosphatidylinositol-calcium second messenger system.,induction:By granulocyte macrophage colony-stimulating factor (GM-CSF), interleukin-5 and n-butyrate.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in the placenta, lung, left and right heart ventricles, heart atrium, leukocytes and differentiated HL-60 granulocytes.,

研究领域 (Research Area)

Calcium;Neuroactive ligand-receptor interaction;

图片 (Image Data)



Immunofluorescence analysis of LOVO cells, using PTAFR Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .