

产品名称: PAI-1 Rabbit Polyclonal Antibody
产品货号: APRab15700



产品概述 (Summary)

产品名称 (Production Name)	PAI-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SERPINE1
别名 (Alternative Names)	SERPINE1; PAI1; PLANH1; Plasminogen activator inhibitor 1; PAI; PAI-1; Endothelial plasminogen activator inhibitor; Serpin E1
基因 ID (Gene ID)	5054.0
蛋白 ID (SwissProt ID)	P05121.The antiserum was produced against synthesized peptide derived from human PAI-1. AA range:266-315

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	47kDa

产品名称: PAI-1 Rabbit Polyclonal Antibody
产品货号: APRab15700




研究背景 (Background)

This gene encodes a member of the serine proteinase inhibitor (serpin) superfamily. This member is the principal inhibitor of tissue plasminogen activator (tPA) and urokinase (uPA), and hence is an inhibitor of fibrinolysis. Defects in this gene are the cause of plasminogen activator inhibitor-1 deficiency (PAI-1 deficiency), and high concentrations of the gene product are associated with thrombophilia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009],disease:Defects in SERPINE1 are the cause of plasminogen activator inhibitor-1 deficiency (PAI-1 deficiency) [MIM:173360]. This deficiency is characterized by abnormal bleeding due to SERPINE1 defect in the plasma.,disease:High concentrations of SERPINE1 have been associated with thrombophilia [MIM:188050]; an autosomal dominant disorder in which affected individuals are prone to develop serious spontaneous thrombosis.,function:This inhibitor acts as 'bait' for tissue plasminogen activator, urokinase, and protein C. Its rapid interaction with TPA may function as a major control point in the regulation of fibrinolysis.,online information:Plasminogen activator inhibitor-1 entry,PTM:Inactivated by proteolytic attack of the urokinase-type (u-PA) and the tissue-type (TPA), cleaving the 369-Arg|-Met-370 bond.,similarity:Belongs to the serpin family.,subunit:Interacts with VTN. Binds LRP1B; binding is followed by internalization and degradation.,tissue specificity:Found in plasma and platelets and in endothelial, hepatoma and fibrosarcoma cells,

研究领域 (Research Area)


p53;Complement and coagulation cascades;

图片 (Image Data)




Immunohistochemistry analysis of PAI-1 antibody in paraffin-embedded human brain tissue.


产品名称: PAI-1 Rabbit Polyclonal Antibody
产品货号: APRab15700



Western blot analysis of lysate from Jurkat cells, using PAI-1 antibody.



Western Blot analysis of various cells using PAI-1 Polyclonal Antibody



Western Blot analysis of Jurkat cells using PAI-1 Polyclonal Antibody

注意事项 (Note)

For research use only .