产品名称: P-cadherin Rabbit Polyclonal Antibody

产品货号: APRab15813



产品概述 (Summary)

产品名称 (Production Name) P-cadherin Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CDH3

别名 (Alternative Names) CDH3; CDHP; Cadherin-3; Placental cadherin; P-cadherin

基因 ID (Gene ID) 1001.0

P22223.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human CDH3. AA range:51-100

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 86kDa

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: P-cadherin Rabbit Polyclonal Antibody

产品货号: APRab15813




This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a gene cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene are associated with hypotrichosis with juvenile macular dystrophy and ectodermal dysplasia, ectrodactyly, and macular dystrophy syndrome (EEMS). [provided by RefSeq, Nov 2015], disease: Defects in CDH3 are the cause of ectodermal dysplasia with ectrodactyly and macular dystrophy (EEM) [MIM:225280]; also known as EEM syndrome, Albrectsen-Svendsen syndrome or Ohdo-Hirayama-Terawaki syndrome. Ectodermal dysplasia defines a heterogeneous group of disorders due to abnormal development of two or more ectodermal structures. EEM is an autosomal recessive condition characterized by features of ectodermal dysplasia such as sparse eyebrows and scalp hair, and selective tooth agenesis associated with macular dystrophy and ectrodactyly, disease: Defects in CDH3 are the cause of hypotrichosis with juvenile macular dystrophy (HJMD) [MIM:601553]. HJMD is a rare autosomal recessive disorder characterized by early hair loss heralding severe degenerative changes of the retinal macula and culminating in blindness during the second to third decade of life., function: Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types., online information: Retina International's Scientific Newsletter, similarity: Contains 5 cadherin domains., subunit: Interacts with CDCP1., tissue specificity:Expressed in some normal epithelial tissues and in some carcinoma cell lines.,

研究领域 (Research Area)

Cell adhesion molecules (CAMs);

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using CDH3 Antibody. The picture on the right is blocked with the synthesized peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838


产品名称: P-cadherin Rabbit Polyclonal Antibody

产品货号: APRab15813






Western blot analysis of lysates from K562 cells, using CDH3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using Cytochrome P450 2C19 antibody.



Western Blot analysis of K562 cells using P-cadherin Polyclonal Antibody

注意事项 (Note)

For research use only.