

产品名称: PC-PLD1 Rabbit Polyclonal Antibody
产品货号: APRab15861



产品概述 (Summary)

产品名称 (Production Name)	PC-PLD1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PLD1
别名 (Alternative Names)	PLD1; Phospholipase D1; PLD 1; hPLD1; Choline phosphatase 1; Phosphatidylcholine-hydrolyzing phospholipase D1
基因 ID (Gene ID)	5337.0
蛋白 ID (SwissProt ID)	Q13393. The antiserum was produced against synthesized peptide derived from human PLD1. AA range: 527-576

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	

产品名称: PC-PLD1 Rabbit Polyclonal Antibody
产品货号: AP Rab15861




研究背景 (Background)

This gene encodes a phosphatidylcholine-specific phospholipase which catalyzes the hydrolysis of phosphatidylcholine in order to yield phosphatidic acid and choline. The enzyme may play a role in signal transduction and subcellular trafficking. Alternative splicing results in multiple transcript variants with both catalytic and regulatory properties. [provided by RefSeq, Sep 2011], catalytic activity: A phosphatidylcholine + H₂O = choline + a phosphatidate., enzyme regulation: Stimulated by phosphatidylinositol 4,5-bisphosphate and phosphatidylinositol 3,4,5-trisphosphate, activated by the phosphokinase C- α , by the ADP-ribosylation factor-1 (ARF-1), and to a lesser extent by GTP-binding proteins: RHO A, RAC-1 and CDC42. Inhibited by oleate., function: Implicated as a critical step in numerous cellular pathways, including signal transduction, membrane trafficking, and the regulation of mitosis. May be involved in the regulation of perinuclear intravesicular membrane traffic., online information: Phospholipase D entry, similarity: Belongs to the phospholipase D family., similarity: Contains 1 PH domain., similarity: Contains 1 PX (phox homology) domain., similarity: Contains 2 PLD phosphodiesterase domains., subunit: Interacts with PIP5K1A., tissue specificity: Expressed abundantly in the pancreas and heart and at high levels in brain, placenta, spleen, uterus and small intestine.,

研究领域 (Research Area)

Glycerophospholipid metabolism; Ether lipid metabolism; Endocytosis; Fc gamma R-mediated phagocytosis; GnRH; Pathways in cancer; Pancreatic cancer;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using PLD1 Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .