

产品名称: Peroxin 19 Rabbit Polyclonal Antibody
产品货号: AP Rab15985



产品概述 (Summary)

产品名称 (Production Name)	Peroxin 19 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PEX19
别名 (Alternative Names)	PEX19; HK33; PXF; OK/SW-cl.22; Peroxisomal biogenesis factor 19; 33 kDa housekeeping protein; Peroxin-19; Peroxisomal farnesylated protein
基因 ID (Gene ID)	5824.0
蛋白 ID (SwissProt ID)	P40855.The antiserum was produced against synthesized peptide derived from human PEX19. AA range:219-268

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	33kDa

产品名称: Peroxin 19 Rabbit Polyclonal Antibody
产品货号: AP Rab15985



研究背景 (Background)


peroxisomal biogenesis factor 19 (PEX19) Homo sapiens This gene is necessary for early peroxisomal biogenesis. It acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. These disorders have at least 14 complementation groups, with more than one phenotype being observed for some complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of Zellweger syndrome (ZWS), as well as peroxisome biogenesis disorder complementation group 14 (PBD-CG14), which is an alternative product. Experimental confirmation may be lacking for some isoforms. Disease: Defects in PEX19 are a cause of Zellweger syndrome (ZWS) [MIM:214100]. ZWS is a fatal peroxisome biogenesis disorder characterized by dysmorphic facial features, hepatomegaly, ocular abnormalities, renal cysts, hearing impairment, profound psychomotor retardation, severe hypotonia and neonatal seizures. Death occurs within the first year of life. Disease: Defects in PEX19 are the cause of peroxisome biogenesis disorder complementation group 14 (PBD-CG14) [MIM:600279]; also known as PBD-CGJ. PBD refers to a group of peroxisomal disorders arising from a failure of protein import into the peroxisomal membrane or matrix. The PBD group is comprised of four disorders: Zellweger syndrome (ZWS), neonatal adrenoleukodystrophy (NALD), infantile Refsum disease (IRD), and classical rhizomelic chondrodysplasia punctata (RCDP). ZWS, NALD and IRD are distinct from RCDP and constitute a clinical continuum of overlapping phenotypes known as the Zellweger spectrum. The PBD group is genetically heterogeneous with at least 14 distinct genetic groups as concluded from complementation studies. Function: Necessary for early peroxisomal biogenesis. Acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Binds and stabilizes newly synthesized PMPs in the cytoplasm by interacting with their hydrophobic membrane-spanning domains, and targets them to the peroxisome membrane by binding to the integral membrane protein PEX3. Excludes CDKN2A from the nucleus and prevents its interaction with MDM2, which results in active degradation of TP53. Similarity: Belongs to the peroxin-19 family. Subcellular location: Mainly cytoplasmic. Some fraction membrane-associated to the outer surface of peroxisomes. Subunit: Interacts with a broad range of peroxisomal membrane proteins, including PEX3, PEX10, PEX11A, PEX11B, PEX12, PEX13, PEX14 and PEX16, PXMP2/PMP22, PXMP4/PMP24, SLC25A17/PMP34, ABCD1/ALDP, ABCD2/ALDRP, and ABCD3/PMP70. Also interacts with the tumor suppressor CDKN2A/p19ARF. Tissue specificity: Ubiquitously expressed. Isoform 1 is strongly predominant in all tissues except in utero where isoform 2 is the main form.

研究领域 (Research Area)


Tags & Cell Markers; Subcellular Markers; Organelles; Peroxisome; Signal Transduction; Protein Trafficking; Organelle Proteins

图片 (Image Data)

产品名称: Peroxin 19 Rabbit Polyclonal Antibody
产品货号: AP Rab15985



Western blot analysis of lysates from MCF-7 cells, using PEX19 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using PEX19 antibody.

注意事项 (Note)

For research use only .