

产品名称: PHLDA3 Rabbit Polyclonal Antibody
产品货号: APRab16078



产品概述 (Summary)

产品名称 (Production Name)	PHLDA3 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PHLDA3
别名 (Alternative Names)	PHLDA3; TIH1; Pleckstrin homology-like domain family A member 3; TDAG51/lpl homolog 1
基因 ID (Gene ID)	23612.0
蛋白 ID (SwissProt ID)	Q9Y5J5.The antiserum was produced against synthesized peptide derived from human PHLDA3. AA range:21-70

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	15kDa

产品名称: PHLDA3 Rabbit Polyclonal Antibody
产品货号: AP Rab16078




研究背景 (Background)

domain:The PH domain binds phosphoinositides with a broad specificity. It competes with the PH domain of AKT1 and directly interferes with AKT1 binding to phosphatidylinositol 4,5-bisphosphate (PIP2) and phosphatidylinositol 3,4,5-triphosphate (PIP3), preventing AKT1 association to membrane lipids and subsequent activation of AKT1 signaling.,function:p53/TP53-regulated repressor of Akt/AKT1 signaling. Represses AKT1 by preventing AKT1-binding to membrane lipids, thereby inhibiting AKT1 translocation to the cellular membrane and activation. Contributes to p53/TP53-dependent apoptosis by repressing AKT1 activity. Its directs transcription regulation by p53/TP53 may explain how p53/TP53 can negatively regulate AKT1. May acts as a tumor suppressor.,induction:By p53/TP53; expression is directly activated by p53/TP53. p53/TP53 phosphorylation on 'Ser-15' is required to activate the PHLDA3 promoter.,miscellaneous:PHLDA3 genomic locus is frequently observed in primary lung cancers, suggesting a role in tumor suppression.,similarity:Belongs to the PHLDA3 family.,similarity:Contains 1 PH domain.,tissue specificity:Widely expressed with lowest expression in liver and spleen.,domain:The PH domain binds phosphoinositides with a broad specificity. It competes with the PH domain of AKT1 and directly interferes with AKT1 binding to phosphatidylinositol 4,5-bisphosphate (PIP2) and phosphatidylinositol 3,4,5-triphosphate (PIP3), preventing AKT1 association to membrane lipids and subsequent activation of AKT1 signaling.,function:p53/TP53-regulated repressor of Akt/AKT1 signaling. Represses AKT1 by preventing AKT1-binding to membrane lipids, thereby inhibiting AKT1 translocation to the cellular membrane and activation. Contributes to p53/TP53-dependent apoptosis by repressing AKT1 activity. Its directs transcription regulation by p53/TP53 may explain how p53/TP53 can negatively regulate AKT1. May acts as a tumor suppressor.,induction:By p53/TP53; expression is directly activated by p53/TP53. p53/TP53 phosphorylation on 'Ser-15' is required to activate the PHLDA3 promoter.,miscellaneous:PHLDA3 genomic locus is frequently observed in primary lung cancers, suggesting a role in tumor suppression.,similarity:Belongs to the PHLDA3 family.,similarity:Contains 1 PH domain.,tissue specificity:Widely expressed with lowest expression in liver and spleen.,

研究领域 (Research Area)

Cell Biology; Other Antibodies

图片 (Image Data)



产品名称: PHLDA3 Rabbit Polyclonal Antibody
产品货号: APRab16078



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PHLDA3 Antibody. The picture on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using PHLDA3 Polyclonal Antibody

注意事项 (Note)

For research use only .