

产品名称: Plk Rabbit Polyclonal Antibody
产品货号: APRab16273



产品概述 (Summary)

产品名称 (Production Name)	Plk Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PLK1
别名 (Alternative Names)	PLK1; PLK; Serine/threonine-protein kinase PLK1; Polo-like kinase 1; PLK-1; Serine/threonine-protein kinase 13; STPK13
基因 ID (Gene ID)	5347.0
蛋白 ID (SwissProt ID)	P53350.The antiserum was produced against synthesized peptide derived from human PLK1. AA range:176-225

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	70kDa

产品名称: Plk Rabbit Polyclonal Antibody
产品货号: AP Rab16273


研究背景 (Background)

The Ser/Thr protein kinase encoded by this gene belongs to the CDC5/Polo subfamily. It is highly expressed during mitosis and elevated levels are found in many different types of cancer. Depletion of this protein in cancer cells dramatically inhibited cell proliferation and induced apoptosis; hence, it is a target for cancer therapy. [provided by RefSeq, Sep 2015], catalytic activity: ATP + a protein = ADP + a phosphoprotein., developmental stage: Accumulates to a maximum during the G2 and M phases, declines to a nearly undetectable level following mitosis and throughout G1 phase, and then begins to accumulate again during S phase., enzyme regulation: Activated by serine and threonine phosphorylation., function: Serine/threonine-protein kinase that performs several important functions throughout M phase of the cell cycle, including the regulation of centrosome maturation and spindle assembly, the removal of cohesins from chromosome arms, the inactivation of APC/C inhibitors, and the regulation of mitotic exit and cytokinesis., induction: By growth-stimulating agents., PTM: Autophosphorylation and phosphorylation of Ser-137 are not significant events during activation of PLK1 in M phase., PTM: Catalytic activity is enhanced by phosphorylation of Thr-210 and/or Ser-137., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. CDC5/Polo subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 2 POLO box domains., subunit: Interacts with CEP170 and EVI5. Interacts and phosphorylates ERCC6L. Interacts with FAM29A., tissue specificity: Placenta and colon.,

研究领域 (Research Area)


Cell_Cycle_G1S; Cell_Cycle_G2M_DNA; Oocyte meiosis; Progesterone-mediated oocyte maturation;

图片 (Image Data)



Immunofluorescence analysis of A549 cells, using PLK1 Antibody. The picture on the right is blocked with the synthesized peptide.

产品名称: Plk Rabbit Polyclonal Antibody
产品货号: APRab16273



Western blot analysis of lysates from LOVO cells, treated with PMA 125ng/ml 30 ' , using PLK1 Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .