

产品名称: PRC1 Rabbit Polyclonal Antibody
产品货号: AP Rab16463



产品概述 (Summary)

产品名称 (Production Name)	PRC1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PRC1
别名 (Alternative Names)	Protein regulator of cytokinesis 1
基因 ID (Gene ID)	9055.0
蛋白 ID (SwissProt ID)	O43663.Synthetic peptide from human protein at AA range: 460-520

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	72kDa

研究背景 (Background)

This gene encodes a protein that is involved in cytokinesis. The protein is present at high levels during the S and G2/M

产品名称: PRC1 Rabbit Polyclonal Antibody
产品货号: AP Rab16463




phases of mitosis but its levels drop dramatically when the cell exits mitosis and enters the G1 phase. It is located in the nucleus during interphase, becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. This protein has been shown to be a substrate of several cyclin-dependent kinases (CDKs). It is necessary for polarizing parallel microtubules and concentrating the factors responsible for contractile ring assembly. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012],function:KIF4A translocates PRC1 to the plus ends of interdigitating spindle microtubules during the metaphase to anaphase transition, an essential step for the formation of an organized central spindle midzone and midbody and for successful cytokinesis. Required for KIF14 localization to the central spindle and midbody. Acts as a microtubule-binding and bundling protein both in vivo and vitro. May function as an in vivo cyclin-CDK substrate.,PTM:Phosphorylated; very weak in G1/S phase cells. Much higher levels of phosphorylation are detected at later cell cycle phases, reaching a maximum during mitosis.,similarity:Belongs to the MAP65/ASE1 family.,subcellular location:Predominantly localized to the nucleus of interphase cells. During mitosis becomes associated with the mitotic spindle poles and localizes with the cell midbody during cytokinesis.,subunit:Interacts with the C-terminal Rho-GAP domain and the basic region of RACGAP1. The interaction with RACGAP1 inhibits its GAP activity towards Cdc42 in vitro, which may be required for maintaining normal spindle morphology. Interacts separately via its N-terminal region with the C-terminus of CENPE, KIF4A and KIF23 during late mitosis. Interacts with KIF14 and KIF20A.,

研究领域 (Research Area)


Cell Biology

图片 (Image Data)



Western blot analysis of lysates from HeLa cells, using PRC1 antibody.

产品名称: PRC1 Rabbit Polyclonal Antibody
产品货号: APRab16463



Western blot analysis of 293T HeLa 3T3 lysate, antibody was diluted at 500. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .