

产品名称: Puratrophin 1 Rabbit Polyclonal Antibody
产品货号: AP Rab16698



产品概述 (Summary)

产品名称 (Production Name)	Puratrophin 1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PLEKHG4 PLEKHG4; PRTPHN1; Puratrophin-1; Pleckstrin homology domain-containing
别名 (Alternative Names)	family G member 4; PH domain-containing family G member 4; Purkinje cell atrophy-associated protein 1
基因 ID (Gene ID)	25894.0
蛋白 ID (SwissProt ID)	Q58EX7.The antiserum was produced against synthesized peptide derived from human PLEKHG4. AA range:654-703

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	135kDa

产品名称: Puratrophin 1 Rabbit Polyclonal Antibody
产品货号: AP Rab16698




研究背景 (Background)

The protein encoded by this gene can function as a guanine nucleotide exchange factor (GEF) and may play a role in intracellular signaling and cytoskeleton dynamics at the Golgi apparatus. Polymorphisms in the region of this gene have been found to be associated with spinocerebellar ataxia in some study populations. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015], disease: Defects in PLEKHG4 are the cause of spinocerebellar ataxia 16q22-linked (SCA16q22) [MIM:117210]; also known as pure spinocerebellar ataxia Japanese type or SCA4 pure Japanese type. Spinocerebellar ataxia is a clinically and genetically heterogeneous group of cerebellar disorders. Patients show progressive incoordination of gait and often poor coordination of hands, speech and eye movements, due to degeneration of the cerebellum with variable involvement of the brainstem and spinal cord. SCA16q22 belongs to the autosomal dominant cerebellar ataxias type III (ADCA III) which are characterized by pure cerebellar ataxia without additional signs., function: Possible role in intracellular signaling and cytoskeleton dynamics at the Golgi., similarity: Contains 1 DH (DBL-homology) domain., similarity: Contains 1 PH domain., tissue specificity: Expressed in kidney, Leydig cells in the testis, epithelial cells in the prostate gland and Langerhans islet in the pancreas. Isoform 1 and isoform 3 are strongly expressed in Purkinje cells and to a lower extent in other neurons (at protein level). Widely expressed at low levels. More strongly expressed in testis and pancreas.,


研究领域 (Research Area)

Neuroscience; Neurology process; Neurodegenerative disease

图片 (Image Data)



产品名称: Puratrophin 1 Rabbit Polyclonal Antibody
产品货号: APRab16698



Western blot analysis of the lysates from HUVEC cells using PLEKHG4 antibody.

注意事项 (Note)

For research use only .