

产品名称: Rac1/2/3/CDC42 Rabbit Polyclonal Antibody
产品货号: AP Rab16822



产品概述 (Summary)

产品名称 (Production Name)	Rac1/2/3/CDC42 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	RAC3
别名 (Alternative Names)	RAC1; TC25; MIG5; Ras-related C3 botulinum toxin substrate 1; Cell migration-inducing gene 5 protein; Ras-like protein TC25; p21-Rac1; RAC2; Ras-related C3 botulinum toxin substrate 2; GX; Small G protein; p21-Rac2; RAC3; Ras-related C3 bot
基因 ID (Gene ID)	5879/5880/5881/998
蛋白 ID (SwissProt ID)	P63000/P15153/P60763/P60953.The antiserum was produced against synthesized peptide derived from human Rac1/CDC42. AA range:38-87

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
----------------------	---

产品名称: Rac1/2/3/CDC42 Rabbit Polyclonal Antibody
产品货号: AP Rab16822



蛋白分子量 (Molecular Weight) 26kDa

研究背景 (Background)

The protein encoded by this gene is a GTPase which belongs to the RAS superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization, and the activation of protein kinases. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009], domain: The effector region mediates interaction with DEF6, enzyme regulation: Regulated by guanine nucleotide exchange factors (GEFs) which promote the exchange of bound GDP for free GTP, GTPase activating proteins (GAPs) which increase the GTP hydrolysis activity, and GDP dissociation inhibitors which inhibit the dissociation of the nucleotide from the GTPase, function: Isoform B has an accelerated GEF-independent GDP/GTP exchange and an impaired GTP hydrolysis, which is restored partially by GTPase-activating proteins. It is able to bind to the GTPase-binding domain of PAK but not full-length PAK in a GTP-dependent manner, suggesting that the insertion does not completely abolish effector interaction, function: Plasma membrane-associated small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate cellular responses such as secretory processes, phagocytosis of apoptotic cells, epithelial cell polarization and growth-factor induced formation of membrane ruffles, similarity: Belongs to the small GTPase superfamily. Rho family, subcellular location: Inner surface of plasma membrane possibly with attachment requiring prenylation of the C-terminal cysteine (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV, subunit: Interacts with the GEF proteins PREX1, RASGRF2, DOCK1, DOCK2 and DOCK7, which promote the exchange between GDP and GTP, and therefore activate it. Interacts with PARD6A, PARD6B and PARD6G in a GTP-dependent manner. Part of a quaternary complex containing PARD3, some PARD6 protein (PARD6A, PARD6B or PARD6G) and some atypical PKC protein (PRKCI or PRKCZ), which plays a central role in epithelial cell polarization. Found in a trimeric complex composed of DOCK1 and ELMO1, which plays a central role in phagocytosis of apoptotic cells. Interacts with RALBP1 via its effector domain. Interacts with PLXNB1. Part of a complex with MAP2K3, MAP3K3, CCM2 and DEF6. Interacts with BAIAP2, BAIAP2L1, CYFIP1/SRA-1 and DEF6. Interacts with Y.pseudotuberculosis YPKA and PLCB2. Interacts with NOXA1. Interacts with ARHGEF2. Interacts with NISCH, tissue specificity: Isoform B is predominantly identified in skin and epithelial tissues from the intestinal tract. The expression of isoform B is elevated in colorectal tumors at various stages of neoplastic progression, as compared to their respective adjacent tissues.,


研究领域 (Research Area)

MAPK_ERK_Growth; MAPK_G_Protein; Chemokine; WNT; WNT-T CELL Axon guidance; VEGF; Focal adhesion; Adherens_Junction; Toll_Like; Natural killer cell mediated cytotoxicity; B_Cell_Antigen; Fc epsilon RI; Fc gamma R-mediated phagocytosis; Leukocyte transendothelial migration; Neurotrophin; Regulates Actin and Cytoskeleton; Amyotrophic lateral sclerosis (ALS); Epithelial cell signaling in Helicobacter pylori infection; Pathways in cancer; Colorectal cancer; Renal cell carcinoma; Pancreatic cancer; Viral myocarditis;


产品名称: Rac1/2/3/CDC42 Rabbit Polyclonal Antibody
产品货号: AP Rab16822




图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Rac1/CDC42 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from NIH/3T3 cells, treated with EGF 200ng/ml 30', using Rac1/CDC42 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Rac1/2/3/CDC42 Polyclonal Antibody

产品名称: Rac1/2/3/CDC42 Rabbit Polyclonal Antibody
产品货号: AP Rab16822



Western Blot analysis of 3T3 cells using Rac1/2/3/CDC42 Polyclonal Antibody

注意事项 (Note)

For research use only .