

产品名称: Rad50 Rabbit Polyclonal Antibody
产品货号: APRab16839



产品概述 (Summary)

产品名称 (Production Name)	Rad50 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	RAD50
别名 (Alternative Names)	RAD50; DNA repair protein RAD50; hRAD50
基因 ID (Gene ID)	10111.0
蛋白 ID (SwissProt ID)	Q92878.The antiserum was produced against synthesized peptide derived from human RAD50. AA range:681-730

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	153kDa

研究背景 (Background)

产品名称: Rad50 Rabbit Polyclonal Antibody
产品货号: AP Rab16839




RAD50 double strand break repair protein(RAD50) Homo sapiens The protein encoded by this gene is highly similar to Saccharomyces cerevisiae Rad50, a protein involved in DNA double-strand break repair. This protein forms a complex with MRE11 and NBS1. The protein complex binds to DNA and displays numerous enzymatic activities that are required for nonhomologous joining of DNA ends. This protein, cooperating with its partners, is important for DNA double-strand break repair, cell cycle checkpoint activation, telomere maintenance, and meiotic recombination. Knockout studies of the mouse homolog suggest this gene is essential for cell growth and viability. Mutations in this gene are the cause of Nijmegen breakage syndrome-like disorder.[provided by RefSeq, Apr 2010],cofactor: Binds 1 zinc ion per homodimer.,domain: The zinc-hook, which separates the large intramolecular coiled coil regions, contains 2 Cys residues that coordinate one molecule of zinc with the help of the 2 Cys residues of the zinc-hook of another RAD50 molecule, thereby forming a V-shaped homodimer. The two heads of the homodimer, which constitute the ATP-binding domain, interact with the MRE11A homodimer.,function: Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis. The complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11A. RAD50 may be required to bind DNA ends and hold them in close proximity. This could facilitate searches for short or long regions of sequence homology in the recombining DNA templates, and may also stimulate the activity of DNA ligases and/or restrict the nuclease activity of MRE11A to prevent nucleolytic degradation past a given point. The complex may also be required for DNA damage signaling via activation of the ATM kinase. In telomeres the MRN complex may modulate t-loop formation.,miscellaneous: In case of infection by adenovirus E4, the MRN complex is inactivated and degraded by viral oncoproteins, thereby preventing concatenation of viral genomes in infected cells.,PTM: Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution: Contaminating sequence. Potential poly-A sequence.,similarity: Belongs to the SMC family. RAD50 subfamily.,similarity: Contains 1 zinc-hook domain.,subcellular location: Localizes to discrete nuclear foci after treatment with genotoxic agents.,subunit: Component of the MRN complex composed of two heterodimers RAD50/MRE11A associated with a single NBN. Component of the BASC complex, at least composed of BRCA1, MSH2, MSH6, MLH1, ATM, BLM, RAD50, MRE11A and NBN. Found in a complex with TERF2. Interacts with RINT1. Interacts with BRCA1 via its N-terminal domain. Interacts with DCLRE1C/Artemis.,tissue specificity: Expressed at very low level in most tissues, except in testis where it is expressed at higher level. Expressed in fibroblasts.,

研究领域 (Research Area)


Homologous recombination;Non-homologous end-joining;

图片 (Image Data)

产品名称: Rad50 Rabbit Polyclonal Antibody
产品货号: APRab16839



Western blot analysis of lysates from COLO205, 293, and HUVEC cells, using RAD50 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Rad50 Polyclonal Antibody diluted at 1: 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA) .

注意事项 (Note)

For research use only .