

产品名称: RGS10 Rabbit Polyclonal Antibody
产品货号: APRab17089



产品概述 (Summary)

产品名称 (Production Name)	RGS10 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	RGS10
别名 (Alternative Names)	RGS10; Regulator of G-protein signaling 10; RGS10
基因 ID (Gene ID)	6001.0
蛋白 ID (SwissProt ID)	O43665.The antiserum was produced against synthesized peptide derived from human RGS10. AA range:80-129

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	20kDa

研究背景 (Background)

产品名称: RGS10 Rabbit Polyclonal Antibody
产品货号: AP Rab17089




Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein signaling 10 belongs to this family. All RGS proteins share a conserved 120-amino acid sequence termed the RGS domain. This protein associates specifically with the activated forms of the two related G-protein subunits, G-alpha_{i3} and G-alpha_z but fails to interact with the structurally and functionally distinct G-alpha subunits. Regulator of G protein signaling 10 protein is localized in the nucleus. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Associates specifically with the activated forms of the G protein subunits G(i)-alpha and G(z)-alpha but fails to interact with the structurally and functionally distinct G(s)-alpha subunit. Activity on G(z)-alpha is inhibited by palmitoylation of the G-protein.,PTM:Isoform 3 is phosphorylated on Ser-16.,similarity:Contains 1 RGS domain.,

研究领域 (Research Area)

Signal Transduction; Signaling Pathway; G Protein Signaling; Heterotrimeric G Proteins; Regulators

图片 (Image Data)



Western blot analysis of RGS10 Antibody. The lane on the right is blocked with the RGS10 peptide.



Western Blot analysis of various cells using RGS10 Polyclonal Antibody

产品名称: RGS10 Rabbit Polyclonal Antibody
产品货号: APRab17089



注意事项 (Note)

For research use only .