



产品概述 (Summary)

产品名称 (Production Name) RNase III Drosha Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) DROSHA

DROSHA; RN3; RNASE3L; RNASEN; Ribonuclease 3; Protein Drosha; **别名 (Alternative Names)**

Ribonuclease III; RNase III; p241

基因 ID (Gene ID) 29102.0

Q9NRR4.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human RNase III Drosha. AA range:774-823

产品应用 (Application)

稀释比(Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight) 160kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838






研究背景 (Background)

drosha ribonuclease III(DROSHA) Homo sapiens This gene encodes a ribonuclease (RNase) III double-stranded RNA-specific ribonuclease and subunit of the microprocessor protein complex, which catalyzes the initial processing step of microRNA (miRNA) synthesis. The encoded protein cleaves the stem loop structure from the primary microRNA (primiRNA) in the nucleus, yielding the precursor miRNA (pre-miRNA), which is then exported to the cytoplasm for further processing. In a human cell line lacking a functional copy of this gene, canonical miRNA synthesis is reduced. Somatic mutations in this gene have been observed in human patients with kidney cancer. [provided by RefSeq, Sep 2016],catalytic activity:Endonucleolytic cleavage to 5'-phosphomonoester.,cofactor:Magnesium or manganese.,function:Executes the initial step of microRNA (miRNA) processing in the nucleus, that is cleavage of pri-miRNA to release pre-miRNA. Involved in pre-rRNA processing. Cleaves double-strand RNA and does not cleave single-strand RNA.,online information:The dark side of RNA -Issue 87 of October 2007,similarity:Contains 1 DRBM (double-stranded RNA-binding) domain.,similarity:Contains 2 RNase III domains.,subcellular location:A fraction is translocated to the nucleolus during the S phase of the cell cycle.,subunit:Interacts with Sp1.,tissue specificity:Ubiquitous.,

研究领域 (Research Area)

Epigenetics and Nuclear Signaling; DNA / RNA; RNA Processing RNAi; Dicer; Nuclear Signaling Pathways; Nuclear Receptors; Nuclear Pore Complex


图片 (Image Data)



Western blot analysis of lysates from brain tissue, using RNase III Drosha antibody.


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838






Western blot analysis of SH-SY5Y K562 HELA using RNase III Drosha antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-

003,Inventbiotech,MN,USA)




Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200





Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200

注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838