

产品名称: Sck Rabbit Polyclonal Antibody
产品货号: AP Rab17647



产品概述 (Summary)

产品名称 (Production Name)	Sck Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SHC2 SHC2; SCK; SHCB; SHC-transforming protein 2; Protein Sck; SHC-transforming protein B; Src homology 2 domain-containing-transforming protein C2; SH2 domain protein C2
别名 (Alternative Names)	
基因 ID (Gene ID)	25759.0
蛋白 ID (SwissProt ID)	P98077.The antiserum was produced against synthesized peptide derived from human SHC2. AA range:261-310

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300
蛋白分子量 (Molecular Weight)	59kDa

产品名称: Sck Rabbit Polyclonal Antibody
产品货号: APRab17647




研究背景 (Background)

domain:The PID domain mediates binding to the TrkA receptor.,function:Signaling adapter that couples activated growth factor receptors to signaling pathway in neurons. Involved in the signal transduction pathways of neurotrophin-activated Trk receptors in cortical neurons.,miscellaneous:PubMed:15057824 has shown that SHC2 is poorly phosphorylated by the Trk receptors, in opposite to PubMed:12006576.,PTM:Phosphorylated on tyrosines by the Trk receptors.,similarity:Contains 1 PID domain.,similarity:Contains 1 SH2 domain.,subunit:Interacts with the Trk receptors in a phosphotyrosine-dependent manner and MEGF12. Once activated, binds to GRB2.,tissue specificity:Expressed in brain. Expressed at high level in the hypothalamus and at low level in the caudate nucleus.,domain:The PID domain mediates binding to the TrkA receptor.,function:Signaling adapter that couples activated growth factor receptors to signaling pathway in neurons. Involved in the signal transduction pathways of neurotrophin-activated Trk receptors in cortical neurons.,miscellaneous:PubMed:15057824 has shown that SHC2 is poorly phosphorylated by the Trk receptors, in opposite to PubMed:12006576.,PTM:Phosphorylated on tyrosines by the Trk receptors.,similarity:Contains 1 PID domain.,similarity:Contains 1 SH2 domain.,subunit:Interacts with the Trk receptors in a phosphotyrosine-dependent manner and MEGF12. Once activated, binds to GRB2.,tissue specificity:Expressed in brain. Expressed at high level in the hypothalamus and at low level in the caudate nucleus.,

研究领域 (Research Area)

ErbB_HER;Chemokine;VEGF;Focal adhesion;Natural killer cell mediated cytotoxicity;Neurotrophin;Insulin_Receptor;Glioma;Chronic myeloid leukemia;

图片 (Image Data)



Western blot analysis of lysates from Jurkat cells, using SHC2 Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .