

产品名称: Scn4b Rabbit Polyclonal Antibody
产品货号: APRab17655



产品概述 (Summary)

产品名称 (Production Name)	Scn4b Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SCN4B
别名 (Alternative Names)	SCN4B; Sodium channel subunit beta-4
基因 ID (Gene ID)	6330.0
蛋白 ID (SwissProt ID)	Q8IWT1.The antiserum was produced against synthesized peptide derived from human SCN4B. AA range:61-110

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	27kDa

研究背景 (Background)


产品名称: Scn4b Rabbit Polyclonal Antibody
产品货号: AP Rab17655

The protein encoded by this gene is one of several sodium channel beta subunits. These subunits interact with voltage-gated alpha subunits to change sodium channel kinetics. The encoded transmembrane protein forms interchain disulfide bonds with SCN2A. Defects in this gene are a cause of long QT syndrome type 10 (LQT10). Three protein-coding and one non-coding transcript variant have been found for this gene.[provided by RefSeq, Mar 2009],disease:Defects in SCN4B are the cause of long QT syndrome type 10 (LQT10) [MIM:611819]. Long QT syndromes are heart disorders characterized by a prolonged QT interval on the ECG and polymorphic ventricular arrhythmias. They cause syncope and sudden death in response to exercise or emotional stress. They can present with a sentinel event of sudden cardiac death in infancy.,function:Modulates channel gating kinetics. Causes negative shifts in the voltage dependence of activation of certain alpha sodium channels, but does not affect the voltage dependence of inactivation.,PTM:Contains a number of interchain disulfide bonds with SCN2A.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,subunit:The voltage-sensitive sodium channel consists of an ion conducting pore forming alpha-subunit regulated by one or more beta-1, beta-2, beta-3 and/or beta-4 subunits. Beta-1 and beta-3 are non-covalently associated with alpha, while beta-2 and beta-4 are covalently linked by disulfide bonds. Associates with SCN2A.,tissue specificity:Expressed at a high level in dorsal root ganglia, at a lower level in brain, spinal cord, skeletal muscle and heart.,

研究领域 (Research Area)


Neuroscience; Neurology process; Neural Signal Transduction

图片 (Image Data)



Western blot analysis of lysates from HepG2 and Jurkat cells, using SCN4B Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: Scn4b Rabbit Polyclonal Antibody
产品货号: APRab17655



Western blot analysis of the lysates from HeLa cells using SCN4B antibody.



Western Blot analysis of various cells using Scn4b Polyclonal Antibody

注意事项 (Note)

For research use only .