

产品名称: SIRT2 Rabbit Polyclonal Antibody  
产品货号: APRab17917



## 产品概述 (Summary)

|                        |                                  |
|------------------------|----------------------------------|
| 产品名称 (Production Name) | SIRT2 Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody       |
| 宿主 (Host)              | Rabbit                           |
| 应用 (Application)       | WB,IHC,ELISA                     |
| 种属反应性 (Reactivity)     | Human,Mouse,Rat                  |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Unmodified                                                                                        |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                                           |
|------------------------|---------------------------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | SIRT2                                                                                                                     |
| 别名 (Alternative Names) | SIRT2; SIR2L; SIR2L2; NAD-dependent protein deacetylase sirtuin-2; Regulatory protein SIR2 homolog 2; SIR2-like protein 2 |
| 基因 ID (Gene ID)        | 22933.0                                                                                                                   |
| 蛋白 ID (SwissProt ID)   | Q8IXJ6.The antiserum was produced against synthesized peptide derived from human SIRT2. AA range:321-370                  |

## 产品应用 (Application)

|                          |                                                     |
|--------------------------|-----------------------------------------------------|
| 稀释比 (Dilution Ratio)     | WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000 |
| 蛋白分子量 (Molecular Weight) | 43kDa                                               |

产品名称: SIRT2 Rabbit Polyclonal Antibody  
产品货号: APRab17917




## 研究背景 (Background)

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from alternative splicing of this gene. [provided by RefSeq, Jul 2010], catalytic activity: NAD(+) + an acetylprotein = nicotinamide + O-acetyl-ADP-ribose + a protein., cofactor: Binds 1 zinc ion per subunit., developmental stage: Peaks during mitosis. After mitosis, it is probably degraded by the 26S proteasome., enzyme regulation: Inhibited by Sirtinol, A3 and M15 small molecules. Inhibited by nicotinamide., function: NAD-dependent deacetylase, which deacetylates the 'Lys-40' of alpha-tubulin. Involved in the control of mitotic exit in the cell cycle, probably via its role in the regulation of cytoskeleton. Despite some ability to deacetylate histones in vitro, it is unlikely in vivo., PTM: Phosphorylated at the G2/M transition of the cell cycle., similarity: Belongs to the sirtuin family., similarity: Contains 1 deacetylase sirtuin-type domain., subcellular location: Colocalizes with microtubules., subunit: Interacts with HDAC6, suggesting that these proteins belong to a large complex that deacetylate the cytoskeleton., tissue specificity: Widely expressed. Highly expressed in heart, brain and skeletal muscle, while it is weakly expressed in placenta and lung. Down-regulated in many gliomas suggesting that it may act as a tumor suppressor gene in human gliomas possibly through the regulation of microtubule network.,

## 研究领域 (Research Area)


Protein\_Acetylation

## 图片 (Image Data)




Western blot analysis of SIRT2 Antibody. The lane on the right is blocked with the SIRT2 peptide.

产品名称: SIRT2 Rabbit Polyclonal Antibody  
产品货号: AP Rab17917



Western blot analysis of the lysates from RAW264.7 cells using SIRT2 antibody.



Western Blot analysis of 293 cells using SIRT2 Polyclonal Antibody diluted at 1: 1000

### 注意事项 (Note)

For research use only .