

产品名称: SL9C1 Rabbit Polyclonal Antibody
产品货号: APRab17941



产品概述 (Summary)

产品名称 (Production Name)	SL9C1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SLC9C1 SLC9A10
别名 (Alternative Names)	
基因 ID (Gene ID)	285335.0
蛋白 ID (SwissProt ID)	Q4G0N8.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	129kDa

研究背景 (Background)

SLC9A10 is a member of the sodium-hydrogen exchanger (NHE) family (see SLC9A1, MIM 107310) and is required for male fertility and sperm motility (Wang et al., 2003 [PubMed 14634667]).[supplied by OMIM, Apr 2009],domain:The ion


产品名称: SL9C1 Rabbit Polyclonal Antibody
产品货号: APRab17941



transport-like region is related to the membrane segments of voltage-gated ion channels. Its function is unknown.,function:Sperm-specific sodium/hydrogen exchanger involved in intracellular pH regulation of spermatozoa. Required for sperm motility and fertility. Involved in sperm cell hyperactivation, a step needed for sperm motility which is essential late in the preparation of sperm for fertilization. Required for the expression and bicarbonate regulation of the soluble adenylyl cyclase (sAC).,similarity:Belongs to the monovalent cation:proton antiporter 1 (CPA1) transporter (TC 2.A.36) family.,similarity:Contains 1 cyclic nucleotide-binding domain.,subunit:Interacts with soluble adenylyl cyclase (sAC).,

研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only .