

产品名称: SLP-76 Rabbit Polyclonal Antibody
产品货号: APRab17980



产品概述 (Summary)

产品名称 (Production Name)	SLP-76 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	LCP2
别名 (Alternative Names)	LCP2; Lymphocyte cytosolic protein 2; SH2 domain-containing leukocyte protein of 76 kDa; SLP-76 tyrosine phosphoprotein; SLP76
基因 ID (Gene ID)	3937.0
蛋白 ID (SwissProt ID)	Q13094.Synthesized peptide derived from SLP-76 . at AA range: 270-350

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	60kDa

研究背景 (Background)

产品名称: SLP-76 Rabbit Polyclonal Antibody
产品货号: AP Rab17980




SLP-76 was originally identified as a substrate of the ZAP-70 protein tyrosine kinase following T cell receptor (TCR) ligation in the leukemic T cell line Jurkat. The SLP-76 locus has been localized to human chromosome 5q33 and the gene structure has been partially characterized in mice. The human and murine cDNAs both encode 533 amino acid proteins that are 72% identical and comprised of three modular domains. The NH₂-terminus contains an acidic region that includes a PEST domain and several tyrosine residues which are phosphorylated following TCR ligation. SLP-76 also contains a central proline-rich domain and a COOH-terminal SH2 domain. A number of additional proteins have been identified that associate with SLP-76 both constitutively and inducibly following receptor ligation, supporting the notion that SLP-76 functions as an adaptor or scaffold protein. Studies using SLP-76 deficient T cell lines: The SH2 domain mediates interaction with SHB, function: Involved in T-cell antigen receptor mediated signaling, PTM: Phosphorylated after T-cell receptor activation by ZAP-70, similarity: Contains 1 SAM (sterile alpha motif) domain, similarity: Contains 1 SH2 domain, subunit: Interacts with SLA. Interacts with CBLB (By similarity). Interacts with the adapter proteins GRB2 and FYN. Interacts with SHB. Interacts with PRAM1, tissue specificity: Highly expressed in spleen, thymus, and peripheral blood leukocytes. Highly expressed also in T-cell and monocytic cell lines, expressed at lower level in B-cell lines. Not detected in fibroblast or neuroblastoma cell lines.

研究领域 (Research Area)


Natural killer cell mediated cytotoxicity; T_Cell_Receptor; Fc epsilon RI;

图片 (Image Data)



Western Blot analysis of extracts from Jurkat cells, using SLP-76 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

产品名称: SLP-76 Rabbit Polyclonal Antibody
产品货号: APRab17980



Immunohistochemical analysis of paraffin-embedded human-prostate-cancer, antibody was diluted at 1:100

注意事项 (Note)

For research use only .