

产品名称: SMG9 Rabbit Polyclonal Antibody
产品货号: APRab18021



产品概述 (Summary)

产品名称 (Production Name)	SMG9 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SMG9 C19orf61
别名 (Alternative Names)	
基因 ID (Gene ID)	56006.0
蛋白 ID (SwissProt ID)	Q9H0W8.Synthesized peptide derived from human protein . at AA range: 70-150

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	57kDa

研究背景 (Background)

SMG9, nonsense mediated mRNA decay factor(SMG9) Homo sapiens This gene encodes a regulatory subunit of the

产品名称: SMG9 Rabbit Polyclonal Antibody
产品货号: APRab18021




SMG1 complex, which plays a critical role in nonsense-mediated mRNA decay (NMD). Binding of the encoded protein to the SMG1 complex kinase scaffold protein results in the inhibition of its kinase activity. Mutations in this gene cause a multiple congenital anomaly syndrome in human patients, characterized by brain malformation, congenital heart disease and other features. [provided by RefSeq, Jul 2016],function:Component of the SMG1C complex, a mRNA surveillance complex that recognizes and degrades mRNAs containing premature translation termination codons (PTCs) via the nonsense-mediated mRNA decay (NMD). The complex probably acts by associating with ribosomes during translation termination on mRNPs. If an exon junction complex (EJC) is located 50-55 or more nucleotides downstream from the termination codon, smg1 phosphorylates upf1/rent1, triggering nonsense-mediated mRNA decay (NMD). In the SMG1C complex, it is required for the efficient association between smg1 and smg8.,PTM:Phosphorylated by SMG1.,similarity:Belongs to the SMG9 family.,subunit:Component of the SMG1C complex, at least composed of SMG1, SMG8 and SMG9. The SMG1C complex is then recruited on premature translation termination codons (PTCs) to form the ribosome:SURF complex, at least composed of ERF1, ERF3 (ERF3A or ERF3B), EEF2, UPF1/RENT1, SMG1, SMG8 and SMG9.,

研究领域 (Research Area)

Epigenetics and Nuclear Signaling; DNA / RNA; RNA Processing

图片 (Image Data)



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only .