

产品名称: SMRTe Rabbit Polyclonal Antibody

产品货号: APRab18032

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Rat,Mouse
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
分子量(Molecular Weight)	270kDa

抗原信息(Antigen Information)

基因名(Gene Name)	NCOR2 NCOR2; CTG26; Nuclear receptor corepressor 2; N-CoR2; CTG repeat protein 26; SMAP270;
别名(Alternative Names)	Silencing mediator of retinoic acid and thyroid hormone receptor; SMRT; T3 receptor-associating factor; TRAC; Thyroid-; retinoic-acid-receptor-associated
基因 ID(Gene ID)	9612.0
SwissProt ID	Q9Y618
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from human NCOR2. AA range:511-560

研究背景 (Background)

This gene encodes a nuclear receptor co-repressor that mediates transcriptional silencing of certain target genes. The encoded protein is a member of a family of thyroid hormone- and retinoic acid receptor-associated co-repressors. This protein acts as part of a multisubunit complex which includes histone deacetylases to modify chromatin structure that prevents basal transcriptional activity of target genes. Aberrant expression of this gene is associated with certain cancers. Alternate splicing results in multiple transcript variants encoding different isoforms.[provided by RefSeq, Apr 2011],domain:The N-terminal region contains repression functions that are divided into three independent repression domains (RD1, RD2 and RD3). The C-terminal region contains the nuclear receptor-interacting domains that are divided in two separate interaction domains (ID1 and ID2),domain:The two interaction domains (ID) contain a conserved sequence referred to as the CORNR box. This motif is required and sufficient to permit binding to unligated TR and RARS. Sequences flanking the CORNR box determine nuclear hormone receptor specificity.,function:Mediates the transcriptional repression activity of some nuclear receptors by promoting chromatin condensation, thus preventing access of the basal transcription. Isoform 1 and isoform 5 have different affinities for different nuclear receptors.,induction:Regulated during cell cycle progression.,sequence caution:Contaminating sequence. Sequence of unknown origin in the N-terminal part.,sequence caution:Wrong choice of CDS.,similarity:Belongs to the N-CoR nuclear receptor corepressors family.,similarity:Contains 2 SANT domains.,subunit:Interacts with HDAC7 (By similarity). Forms a large corepressor complex that contains SIN3A/B and histone deacetylases HDAC1 and HDAC2. This complex associates with the thyroid (TR) and the retinoid acid receptors (RAR) in the absence of ligand, and may stabilize their interaction with TFIIB. Isoform SRMT interacts with HDAC10. Interacts with MINT. Component of the N-CoR repressor complex, at least composed of NCOR1, NCOR2, HDAC3, TBL1X, TBL1R, CORO2A and GPS2. Interacts with CBFA2T3. Interacts with C1D (By similarity). Interacts with ATXN1L,tissue specificity:Ubiquitous. High levels of expression are detected in lung, spleen and brain.,


研究领域 (Research Area)

Notch;


注意事项 (Note)

For research use only.

图片 (Image Data)



Immunofluorescence analysis of A549 cells, using NCOR2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using NCOR2 Antibody. The picture on the right is blocked with the synthesized peptide.

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>