

产品名称: SODE Rabbit Polyclonal Antibody
产品货号: APRab18100



产品概述 (Summary)

产品名称 (Production Name)	SODE Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SOD3
别名 (Alternative Names)	
基因 ID (Gene ID)	6649.0
蛋白 ID (SwissProt ID)	P08294.Synthesized peptide derived from human SODE AA range: 67-117

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)

This gene encodes a member of the superoxide dismutase (SOD) protein family. SODs are antioxidant enzymes that


产品名称: SODE Rabbit Polyclonal Antibody
产品货号: APRab18100



catalyze the conversion of superoxide radicals into hydrogen peroxide and oxygen, which may protect the brain, lungs, and other tissues from oxidative stress. Proteolytic processing of the encoded protein results in the formation of two distinct homotetramers that differ in their ability to interact with the extracellular matrix (ECM). Homotetramers consisting of the intact protein, or type C subunit, exhibit high affinity for heparin and are anchored to the ECM. Homotetramers consisting of a proteolytically cleaved form of the protein, or type A subunit, exhibit low affinity for heparin and do not interact with the ECM. A mutation in this gene may be associated with increased heart disease risk. [provided by RefSeq, Oct 2015],catalytic activity:2 superoxide + 2 H(+) = O(2) + H(2)O(2),cofactor:Binds 1 copper ion per subunit,cofactor:Binds 1 zinc ion per subunit,function:Destroys radicals which are normally produced within the cells and which are toxic to biological systems,function:Protect the extracellular space from toxic effect of reactive oxygen intermediates by converting superoxide radicals into hydrogen peroxide and oxygen,online information:Superoxide dismutase entry,polymorphism:The variant Gly-231 which is found in about 2.2% of individual displays a 10-fold increased plasma EC-SOD content due to reduced heparin-binding affinity and thus the impairment of its binding ability to endothelial cell surface,similarity:Belongs to the Cu-Zn superoxide dismutase family,subcellular location:99% of EC-SOD is anchored to heparan sulfate proteoglycans in the tissue interstitium, and 1% is located in the vasculature in equilibrium between the plasma and the endothelium,subunit:Homotetramer,tissue specificity:Expressed in blood vessels, heart, lungs, kidney and placenta. Major SOD isoenzyme in extracellular fluids such as plasma, lymph and synovial fluid.,


研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4° overnight) .
2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

产品名称: SODE Rabbit Polyclonal Antibody
产品货号: APRab18100



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4°over night

注意事项 (Note)

For research use only .