

产品名称: SPEG Rabbit Polyclonal Antibody
产品货号: APRab18180



产品概述 (Summary)

产品名称 (Production Name)	SPEG Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SPEG
别名 (Alternative Names)	APEG1 KIAA1297
基因 ID (Gene ID)	10290.0
蛋白 ID (SwissProt ID)	Q15772.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:300, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	359kDa

研究背景 (Background)

This gene encodes a protein with similarity to members of the myosin light chain kinase family. This protein family is required for myocyte cytoskeletal development. Along with the desmin gene, expression of this gene may be controlled by

产品名称: SPEG Rabbit Polyclonal Antibody
产品货号: APRab18180



the desmin locus control region. Mutations in this gene are associated with centronuclear myopathy 5. [provided by RefSeq, Jun 2016],catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Isoform 3 may have a role in regulating the growth and differentiation of arterial smooth muscle cells.,induction:Isoform 3 is quickly down-regulated in response to vascular injury, when ASMC cells change from a quiescent to a proliferative phenotype.,miscellaneous:Expression is under the tight control of the locus control region (LCRs).,PTM:May be autophosphorylated. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 2 fibronectin type-III domains.,similarity:Contains 2 protein kinase domains.,similarity:Contains 9 Ig-like (immunoglobulin-like) domains.,subunit:Isoform 3 is found as a monomer or homodimer.,tissue specificity:Isoform 1 is preferentially expressed in striated muscle. Non-kinase form such as isoform 3 is predominantly expressed in the aorta. Isoform 3 appears to be expressed only in highly differentiated ASMC in normal vessel walls and down-regulated in dedifferentiated ASMC in vivo. In response to vascular injuries ASMC dedifferentiate and change from a quiescent and contractile phenotype to a proliferative and synthetic phenotype. This proliferation of vascular smooth muscle cells is one of the most prominent features of arteriosclerosis.,

研究领域 (Research Area)

Cardiovascular; Vasculature; Vasculature Markers; Arterial

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight.3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only .