

产品名称: SPTN4 Rabbit Polyclonal Antibody
产品货号: APRab18232



产品概述 (Summary)

产品名称 (Production Name)	SPTN4 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SPTBN4
别名 (Alternative Names)	KIAA1642 SPTBN3
基因 ID (Gene ID)	57731.0
蛋白 ID (SwissProt ID)	Q9H254.Synthesized peptide derived from part region of human protein

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:300, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	282kDa

研究背景 (Background)

Spectrin is an actin crosslinking and molecular scaffold protein that links the plasma membrane to the actin cytoskeleton, and functions in the determination of cell shape, arrangement of transmembrane proteins, and organization of organelles.


产品名称: SPTN4 Rabbit Polyclonal Antibody
产品货号: APRab18232



It is composed of two antiparallel dimers of alpha- and beta- subunits. This gene is one member of a family of beta-spectrin genes. The encoded protein localizes to the nuclear matrix, PML nuclear bodies, and cytoplasmic vesicles. A highly similar gene in the mouse is required for localization of specific membrane proteins in polarized regions of neurons. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],similarity:Belongs to the spectrin family.,similarity:Contains 1 PH domain.,similarity:Contains 18 spectrin repeats.,similarity:Contains 2 CH (calponin-homology) domains.,tissue specificity:Abundantly expressed in brain and pancreatic islets.,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only .