

产品名称: SREC Rabbit Polyclonal Antibody
产品货号: APRab18265



产品概述 (Summary)

产品名称 (Production Name)	SREC Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SCARF1 KIAA0149 SREC
别名 (Alternative Names)	
基因 ID (Gene ID)	8578.0
蛋白 ID (SwissProt ID)	Q14162.Synthesized peptide derived from human protein . at AA range: 390-470

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	91kDa

研究背景 (Background)

The protein encoded by this gene is a scavenger receptor that is expressed in endothelial cells. It regulates the uptake of


产品名称: SREC Rabbit Polyclonal Antibody
产品货号: APRab18265



chemically modified low density lipoproteins, including acetylated low density lipoprotein (Ac-LDL), and it may be involved in atherogenesis. This gene is regulated by the transcription factors ZNF444/EZF-2 and SP1. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2013],caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Mediates the binding and degradation of acetylated low density lipoprotein (Ac-LDL). Mediates heterophilic interactions, suggesting a function as adhesion protein.,similarity:Contains 6 EGF-like domains.,subunit:Heterophilic interaction with SREC2 via its extracellular domain. The heterophilic interaction is suppressed by the presence of ligand such as Ac-LDL.,tissue specificity:Endothelial cells.,

研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only .