

产品名称: Synaptotagmin Rabbit Polyclonal Antibody
产品货号: APRab18495



产品概述 (Summary)

产品名称 (Production Name)	Synaptotagmin Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SYT1/SYT2
别名 (Alternative Names)	SYT1; SVP65; SYT; Synaptotagmin-1; Synaptotagmin I; SytI; p65; SYT2; Synaptotagmin-2; Synaptotagmin II; SytII
基因 ID (Gene ID)	6857/127833
蛋白 ID (SwissProt ID)	P21579/Q8N9I0.The antiserum was produced against synthesized peptide derived from human Synaptotagmin. AA range:276-325

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	50kDa

产品名称: Synaptotagmin Rabbit Polyclonal Antibody
产品货号: APRab18495




研究背景 (Background)

The synaptotagmins are integral membrane proteins of synaptic vesicles thought to serve as Ca²⁺ sensors in the process of vesicular trafficking and exocytosis. Calcium binding to synaptotagmin-1 participates in triggering neurotransmitter release at the synapse (Fernandez-Chacon et al., 2001 [PubMed 11242035]).[supplied by OMIM, Jul 2010],cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,domain: The first C2 domain mediates Ca²⁺-dependent phospholipid binding.,domain: The second C2 domain mediates interaction with SV2A and STN2.,function: May have a regulatory role in the membrane interactions during trafficking of synaptic vesicles at the active zone of the synapse. It binds acidic phospholipids with a specificity that requires the presence of both an acidic head group and a diacyl backbone. A Ca²⁺-dependent interaction between synaptotagmin and putative receptors for activated protein kinase C has also been reported. It can bind to at least three additional proteins in a Ca²⁺-independent manner; these are neuexins, syntaxin and AP2.,similarity: Belongs to the synaptotagmin family.,similarity: Contains 2 C2 domains.,subcellular location: Synaptic vesicles and chromaffin granules.,subunit: Homotetramer (Probable). Interacts with SCAMP5, STN2, SV2A, SV2B, SV2C and RIMS1. Forms a complex with SV2B, syntaxin 1 and SNAP25.,

研究领域 (Research Area)


Neuroscience; Neurotransmission; Secretory Vesicles; Munc18; Cell Adhesion Proteins; ECM Proteins; Cell Type Marker; Neuron marker; Synapse marker; Cancer; Cancer Metabolism; Metabolic signaling pathway; Integration of energy metabolism; Metabolism; Pathways and Processes; Metabolic signaling pathways; Energy transfer pathways; Integration of energy; Types of disease; Cancer

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Synaptotagmin Antibody. The picture on the right is blocked with the synthesized peptide.

产品名称: Synaptotagmin Rabbit Polyclonal Antibody
产品货号: APRab18495



Western blot analysis of lysates from 293 cells, treated with Forskolin 40nM 30', using Synaptotagmin Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Synaptotagmin Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only .