

产品名称: Synphilin-1 Rabbit Polyclonal Antibody
产品货号: AP Rab18509



产品概述 (Summary)

产品名称 (Production Name)	Synphilin-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SNCAIP
别名 (Alternative Names)	SNCAIP; Synphilin-1; Sph1; Alpha-synuclein-interacting protein
基因 ID (Gene ID)	9627.0
蛋白 ID (SwissProt ID)	Q9Y6H5.The antiserum was produced against synthesized peptide derived from human Synphilin-1. AA range:797-846

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	100kDa

研究背景 (Background)

产品名称: Synphilin-1 Rabbit Polyclonal Antibody
产品货号: AP Rab18509




This gene encodes a protein containing several protein-protein interaction domains, including ankyrin-like repeats, a coiled-coil domain, and an ATP/GTP-binding motif. The encoded protein interacts with alpha-synuclein in neuronal tissue and may play a role in the formation of cytoplasmic inclusions and neurodegeneration. A mutation in this gene has been associated with Parkinson's disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015], disease: Defects in SNCAIP are a cause of Parkinson disease (PD) [MIM:168600]. PD is a complex, multifactorial disorder that typically manifests after the age of 50 years, although early-onset cases (before 50 years) are known. PD generally arises as a sporadic condition but is occasionally inherited as a simple mendelian trait. Although sporadic and familial PD are very similar, inherited forms of the disease usually begin at earlier ages and are associated with atypical clinical features. PD is characterized by bradykinesia, resting tremor, muscular rigidity and postural instability, as well as by a clinically significant response to treatment with levodopa. The pathology involves the loss of dopaminergic neurons in the substantia nigra and the presence of Lewy bodies (intraneuronal accumulations of aggregated proteins), in surviving neurons in various areas of the brain., miscellaneous: Constructs encoding portions of SNCA and SNCAIP co-transfected in mammalian cells promote cytosolic inclusions resembling the Lewy bodies of Parkinson disease. Coexpression of SNCA, SNCAIP, and PARK2 result in the formation of Lewy body-like ubiquitin-positive cytosolic inclusions. Familial mutations in PARK2 disrupt the ubiquitination of SNCAIP and the formation of the ubiquitin-positive inclusions. These results provide a molecular basis for the ubiquitination of Lewy body-associated proteins and link PARK2 and SNCA in a common pathogenic mechanism through their interaction with SNCAIP., PTM: Ubiquitinated; mediated by SIAH1 or RNF19A and leading to its subsequent proteasomal degradation., similarity: Contains 6 ANK repeats., subunit: Associates with SNCA, RNF19A AND PARK2., tissue specificity: Widely expressed, with highest levels in brain, heart and placenta.,

研究领域 (Research Area)


Parkinson's disease;

图片 (Image Data)




Immunohistochemistry analysis of Synphilin-1 antibody in paraffin-embedded lung carcinoma. tissue.


产品名称: Synphilin-1 Rabbit Polyclonal Antibody
产品货号: APRab18509



Western blot analysis of lysate from COLO205, using Synphilin-1 antibody.



Western Blot analysis of various cells using Synphilin-1 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100 (4°, overnight) .
High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

注意事项 (Note)

For research use only .