产品名称: T2R41 Rabbit Polyclonal Antibody

产品货号: APRab18568



产品概述 (Summary)

产品名称 (Production Name) T2R41 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,ELISA

种属反应性 (Reactivity) Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) TAS2R41

别名 (Alternative Names)

基因 ID (Gene ID) 259287.0

P59536.Synthesized peptide derived from human protein . at AA range: 150-蛋白ID (SwissProt ID)

,

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:5000-1:20000

230

蛋白分子量 (Molecular Weight) 33kDa

研究背景 (Background)

function: Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab18568




the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.,function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.,

研究领域 (Research Area)

Taste transduction;

图片 (Image Data)



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838