

产品名称: TACE Rabbit Polyclonal Antibody
产品货号: APRab18602



产品概述 (Summary)

产品名称 (Production Name)	TACE Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ADAM17 ADAM17; CSVP; TACE; Disintegrin and metalloproteinase domain-containing
别名 (Alternative Names)	protein 17; ADAM 17; Snake venom-like protease; TNF-alpha convertase; TNF-alpha-converting enzyme; CD antigen CD156b
基因 ID (Gene ID)	6868.0
蛋白 ID (SwissProt ID)	P78536.The antiserum was produced against synthesized peptide derived from human ADAM 17. AA range:701-750

产品应用 (Application)

稀释比 (Dilution Ratio)	ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	

产品名称: TACE Rabbit Polyclonal Antibody
产品货号: AP Rab18602

研究背景 (Background)


ADAM metallopeptidase domain 17(ADAM17) Homo sapiens This gene encodes a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biologic processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis. The encoded preproprotein is proteolytically processed to generate the mature protease. The encoded protease functions in the ectodomain shedding of tumor necrosis factor-alpha, in which soluble tumor necrosis factor-alpha is released from the membrane-bound precursor. This protease also functions in the processing of numerous other substrates, including cell adhesion proteins, cytokine and growth factor receptors and epidermal growth factor (EGF) receptor ligands. The encoded protein also plays a prominent role in the activation of catalytic activity: Narrow endopeptidase specificity. Cleaves Pro-Leu-Ala-Gln-Ala-[Val-Arg-Ser-Ser-Ser in the membrane-bound, 26-kDa form of tumor necrosis factor alpha (TNF-alpha). Similarly cleaves other membrane-anchored, cell-surface proteins to 'shed' the extracellular domains., cofactor: Binds 1 zinc ion per subunit., domain: Must be membrane anchored to cleave the different substrates. The cytoplasmic domain is not required for this activity. Only the catalytic domain is essential to shed TNF and p75 TNFR., domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme., function: Cleaves the membrane-bound precursor of TNF-alpha to its mature soluble form. Responsible for the proteolytic release of several other cell-surface proteins, including p75 TNF-receptor, interleukin 1 receptor type II, p55 TNF-receptor, transforming growth factor-alpha, L-selectin, growth hormone receptor, MUC1 and the amyloid precursor protein. Also involved in the activation of Notch pathway., induction: In arthritis-affected cartilage., online information: Tumor necrosis factor alpha-converting enzyme entry, PTM: Phosphorylated. Stimulation by growth factor or phorbol 12-myristate 13-acetate induces phosphorylation of Ser-819 but decreases phosphorylation of Ser-791., PTM: The precursor is cleaved by a furin endopeptidase., similarity: Contains 1 disintegrin domain., similarity: Contains 1 peptidase M12B domain., subunit: Interacts with MAD2L1 and MUC1., tissue specificity: Ubiquitously expressed. Expressed at highest levels in adult heart, placenta, skeletal muscle, pancreas, spleen, thymus, prostate, testes, ovary and small intestine, and in fetal brain, lung, liver and kidney.,

研究领域 (Research Area)

Notch; Alzheimer's disease; Epithelial cell signaling in Helicobacter pylori infection;

图片 (Image Data)

产品名称: TACE Rabbit Polyclonal Antibody
产品货号: APRab18602



Immunofluorescence analysis of NIH/3T3 cells, using ADAM 17 Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .