

产品名称: Tubulin α Rabbit Polyclonal Antibody
产品货号: APRab19418



产品概述 (Summary)

产品名称 (Production Name)	Tubulin α Rabbit Polyclonal Antibody
描述 (Description)	Rabbit Polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.03% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TUBA1A/TUBA1B TUBA1A; TUBA3; Tubulin alpha-1A chain; Alpha-tubulin 3; Tubulin B-alpha-1;
别名 (Alternative Names)	Tubulin alpha-3 chain; TUBA1B; Tubulin alpha-1B chain; Alpha-tubulin ubiquitous; Tubulin K-alpha-1; Tubulin alpha-ubiquitous chain
基因 ID (Gene ID)	7846.0
蛋白 ID (SwissProt ID)	Q71U36/P68363. Synthesized peptide derived from the C-terminal region of human Tubulin α .

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000. ELISA: 1:20000.
蛋白分子量 (Molecular Weight)	50kD

产品名称: Tubulin α Rabbit Polyclonal Antibody
产品货号: APRab19418




研究背景 (Background)

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulins. The genes encoding these microtubule constituents belong to the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes, which are highly conserved among species. This gene encodes alpha tubulin and is highly similar to the mouse and rat Tuba1 genes. Northern blotting studies have shown that the gene expression is predominantly found in morphologically differentiated neurologic cells. This gene is one of three alpha-tubulin genes in a cluster on chromosome 12q. disease: Defects in TUBA1A are the cause of lissencephaly type 3 (LIS3) [MIM:611603]. LIS is characterized by a smooth brain surface due to the absence (agyria) or reduction (pachygyria) of surface convolutions. It is often associated with psychomotor retardation and seizures. LIS3 features include agyria or pachygyria or laminar heterotopia, severe mental retardation, motor delay, variable presence of seizures, and abnormalities of corpus callosum, hippocampus, cerebellar vermis and brainstem. function: Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain. PTM: Undergoes a tyrosination/detyrosination cycle, the cyclic removal and re-addition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), respectively. similarity: Belongs to the tubulin family. subunit: Dimer of alpha and beta chains. tissue specificity: Expressed at a high level in fetal brain.

研究领域 (Research Area)

Gap junction; Pathogenic Escherichia coli infection;

图片 (Image Data)



Western Blot analysis of various cells using Tubulin α Polyclonal Antibody. Secondary antibody was diluted at 1:20000

产品名称: Tubulin α Rabbit Polyclonal Antibody
产品货号: APRab19418



注意事项 (Note)

For research use only .