

产品名称: TULA Rabbit Polyclonal Antibody
产品货号: APRab19430



产品概述 (Summary)

产品名称 (Production Name)	TULA Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	UBASH3A UBASH3A; STS2; Ubiquitin-associated and SH3 domain-containing protein A;
别名 (Alternative Names)	Cbl-interacting protein 4; CLIP4; Suppressor of T-cell receptor signaling 2; STS-2; T-cell ubiquitin ligand 1; TULA-1
基因 ID (Gene ID)	53347.0
蛋白 ID (SwissProt ID)	P57075.The antiserum was produced against synthesized peptide derived from human UBASH3A. AA range:311-360

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	75kDa

产品名称: TULA Rabbit Polyclonal Antibody
产品货号: APRab19430




研究背景 (Background)

This gene encodes one of two family members belonging to the T-cell ubiquitin ligand (TULA) family. Both family members can negatively regulate T-cell signaling. This family member can facilitate growth factor withdrawal-induced apoptosis in T cells, which may occur via its interaction with AIF, an apoptosis-inducing factor. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Aug 2011],function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors, EGFR and PDGFRB, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Homodimer or homooligomer. Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,tissue specificity:Highest expression of UBASH3A in tissues belonging to the immune system, including spleen, peripheral blood leukocytes, thymus and bone marrow.,

研究领域 (Research Area)

Cell Biology; Other Antibodies; Other Antibodies

图片 (Image Data)



Western blot analysis of UBASH3A Antibody. The lane on the right is blocked with the UBASH3A peptide.

注意事项 (Note)

For research use only .