

产品名称: WISP-3 Rabbit Polyclonal Antibody
产品货号: APRab19906



产品概述 (Summary)

产品名称 (Production Name)	WISP-3 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	WISP3 CCN6 UNQ462/PRO790/PRO956
别名 (Alternative Names)	WNT1-inducible-signaling pathway protein 3 (WISP-3) (CCN family member 6)
基因 ID (Gene ID)	8838.0
蛋白 ID (SwissProt ID)	O95389.The antiserum was produced against synthesized peptide derived from the N-terminal region of human WISP3. AA range:1-50

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	55kDa

研究背景 (Background)


产品名称: WISP-3 Rabbit Polyclonal Antibody
产品货号: APRab19906

This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene is overexpressed in colon tumors. It may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. Mutations of this gene are associated with progressive pseudorheumatoid dysplasia, an autosomal recessive skeletal disorder, indicating that the gene is essential for normal postnatal skeletal growth and cartilage homeostasis. Multipledisease:Defects in WISP3 are the cause of progressive pseudorheumatoid arthropathy of childhood (PPAC) [MIM:208230]. PPAC is an autosomal recessive disorder characterized by stiffness and swelling of joints, motor weakness and joint contractures. Signs and symptoms of the disease develop typically between three and eight years of age. This progressive disease is a primary disorder of articular cartilage with continued cartilage loss and destructive bone changes with aging.,function:Appears to be required for normal postnatal skeletal growth and cartilage homeostasis.,similarity:Belongs to the CCN family.,similarity:Contains 1 CTCK (C-terminal cystine knot-like) domain.,similarity:Contains 1 IGFBP N-terminal domain.,similarity:Contains 1 TSP type-1 domain.,tissue specificity:Predominant expression in adult kidney and testis and fetal kidney. Weaker expression found in placenta, ovary, prostate and small intestine. Also expressed in skeletally-derived cells such as synoviocytes and articular cartilage chondrocytes.,

研究领域 (Research Area)


Immunology; Secreted Molecules; CCN Family; Stem Cells; Signaling Pathways; Wnt; Secreted; Mesenchymal Stem Cells; Secreted Molecules

图片 (Image Data)



Western blot analysis of Hela Cell Lysate using antibody. Secondary antibody was diluted at 1:20000

产品名称: WISP-3 Rabbit Polyclonal Antibody
产品货号: APRab19906



Immunohistochemical analysis of paraffin-embedded human-liver-cancer, antibody was diluted at 1:200

注意事项 (Note)

For research use only .