

产品名称: ZAP-70 Rabbit Polyclonal Antibody
产品货号: AP Rab20028



产品概述 (Summary)

产品名称 (Production Name)	ZAP-70 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ZAP70
别名 (Alternative Names)	ZAP70; SRK; Tyrosine-protein kinase ZAP-70; 70 kDa zeta-chain associated protein; Syk-related tyrosine kinase
基因 ID (Gene ID)	7535.0
蛋白 ID (SwissProt ID)	P43403.The antiserum was produced against synthesized peptide derived from human ZAP-70. AA range:286-335

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	70kDa

产品名称: ZAP-70 Rabbit Polyclonal Antibody
产品货号: AP Rab20028




研究背景 (Background)

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq, Jul 2008], catalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate}$, disease: Defects in ZAP70 are the cause of selective T-cell defect (STD) [MIM:176947]. STD is an autosomal recessive form of severe combined immunodeficiency characterized by a selective absence of CD8-type T-cells, domain: The SH2 domain binds to the phosphorylated tyrosine-based activation motif (TAM) of CD3Z, function: Plays a role in T-cell development and lymphocyte activation. Essential for TCR-mediated IL-2 production. Isoform 1 induces TCR-mediated signal transduction, isoform 2 does not, online information: ZAP70 mutation db, PTM: Phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation. Tyr-319 phosphorylation is essential for full activity, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. SYK/ZAP-70 subfamily, similarity: Contains 1 protein kinase domain, similarity: Contains 2 SH2 domains, subcellular location: After antigen stimulation, isoform 1 concentrates at the immunological synapse and isoform 2 remains cytoplasmic, subunit: Interacts with SLA2 when it is phosphorylated. Interacts with CD3Z and with phosphorylated NFAM1. Interacts with CBLB (By similarity). Interacts with CBL and SLA when it is phosphorylated. The association with SLA (or SLA2) and CBL probably leads to its destruction. Interacts with SHB. Interacts with DEF6 (By similarity). Interacts with FCRL3, tissue specificity: Expressed in T- and natural killer cells,

研究领域 (Research Area)


Natural killer cell mediated cytotoxicity; T_Cell_Receptor; Primary immunodeficiency;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using ZAP-70 Antibody. The picture on the right is blocked with the synthesized peptide.

产品名称: ZAP-70 Rabbit Polyclonal Antibody
产品货号: APRab20028



Western blot analysis of lysates from Jurkat cells, using ZAP-70 Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .