

产品名称: ZFP91 Rabbit Polyclonal Antibody
产品货号: APRab20092



产品概述 (Summary)

产品名称 (Production Name)	ZFP91 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ZFP91 ZNF757 FKSG11
别名 (Alternative Names)	ZFP91 ZNF757 FKSG11
基因 ID (Gene ID)	80829.0
蛋白 ID (SwissProt ID)	Q96JP5.The antiserum was produced against synthesized peptide derived from the Internal region of human ZFP91. AA range:401-450

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	60kDa

研究背景 (Background)

产品名称: ZFP91 Rabbit Polyclonal Antibody
产品货号: APRab20092




The protein encoded by this gene is a member of the zinc finger family of proteins. The gene product contains C2H2-type domains, which are the classical zinc finger domains found in numerous nucleic acid-binding proteins. This protein functions as a regulator of the non-canonical NF-kappaB pathway in lymphotoxin-beta receptor signaling. Alternative splicing results in multiple transcript variants. A read-through transcript variant composed of ZFP91 and the downstream CNTF gene sequence has been identified, but it is thought to be non-coding. Read-through transcription of ZFP91 and CNTF has also been observed in mouse. A ZFP91-related pseudogene has also been identified on chromosome 2. [provided by RefSeq, Oct 2010],disease:Overexpressed in most acute myelogenous leukemia (AML) cases (27 over 29),,function:CNTF is a survival factor for various neuronal cell types. Seems to prevent the degeneration of motor axons after axotomy,,function:May be involved in transcriptional regulation. May play an important role in cell proliferation and/or anti-apoptosis,,online information:Ciliary neurotrophic factor entry,similarity:Belongs to the CNTF family,,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family,,similarity:Contains 5 C2H2-type zinc fingers,,subunit:Homodimer,,tissue specificity:Expressed ubiquitously, particularly at high level in testis. Isoform 2 is testis specific,,tissue specificity:Nervous system,,

研究领域 (Research Area)

Cytokine-cytokine receptor interaction;Jak_STAT;

图片 (Image Data)



Western Blot analysis of hepg2, NT28 cells using Antibody diluted at 500. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .