

产品名称: ZNRF2 Rabbit Polyclonal Antibody
产品货号: APRab20294



产品概述 (Summary)

产品名称 (Production Name)	ZNRF2 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF, ELISA
种属反应性 (Reactivity)	Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ZNRF2
别名 (Alternative Names)	ZNRF2; RNF202; E3 ubiquitin-protein ligase ZNRF2; Protein Ells2; RING finger protein 202; Zinc/RING finger protein 2
基因 ID (Gene ID)	223082.0
蛋白 ID (SwissProt ID)	Q8NHG8. The antiserum was produced against synthesized peptide derived from human ZNRF2. AA range: 161-210

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:200-1:1000, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

产品名称: ZNRF2 Rabbit Polyclonal Antibody
产品货号: APRab20294


研究背景 (Background)

domain:The RING-type zinc finger domain is required for E3 ligase activity.,function:May play a role in the establishment and maintenance of neuronal transmission and plasticity via its ubiquitin ligase activity. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Present in presynaptic plasma membranes in neurons.,subunit:Interacts with UBE2N.,tissue specificity:Highly expressed in the brain, with higher expression during development than in adult. Expressed also in mammary glands, testis, colon and kidney.,domain:The RING-type zinc finger domain is required for E3 ligase activity.,function:May play a role in the establishment and maintenance of neuronal transmission and plasticity via its ubiquitin ligase activity. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfer the ubiquitin to targeted substrates.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Present in presynaptic plasma membranes in neurons.,subunit:Interacts with UBE2N.,tissue specificity:Highly expressed in the brain, with higher expression during development than in adult. Expressed also in mammary glands, testis, colon and kidney.,

研究领域 (Research Area)

Cell Biology; Proteolysis / Ubiquitin; Proteasome / Ubiquitin; Ubiquitin E3 Enzymes; RING Finger E3 Ligase

图片 (Image Data)



Immunofluorescence analysis of A549 cells, using ZNRF2 Antibody. The picture on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .