

产品名称: Phospho-EGFR (Tyr1092) Rabbit Monoclonal Antibody
产品货号: AMRe01930



产品概述 (Summary)

产品名称 (Production Name)	Phospho-EGFR (Tyr1092) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	EGFR
别名 (Alternative Names)	EGFR; ERBB; ERBB1; HER1; Epidermal growth factor receptor; Proto-oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1
基因 ID (Gene ID)	1956
蛋白 ID (SwissProt ID)	P00533.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000,IHC 1:50-1:100
蛋白分子量 (Molecular Weight)	Calculated MW: 134 kDa; Observed MW: 175 kDa

产品名称: Phospho-EGFR (Tyr1092) Rabbit Monoclonal Antibody
产品货号: AMRe01930




研究背景 (Background)

EGFR is a receptor tyrosine kinase. Receptor for epidermal growth factor (EGF) and related growth factors including TGF- α , amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. . A single-pass transmembrane tyrosine kinase. Ligand binding to this receptor results in receptor dimerization, autophosphorylation (in trans), activation of various downstream signaling molecules and lysosomal degradation.


研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Western blot analysis of EGFR (Phospho-Tyr1092) in C6, 3T3, HeLa lysates using Phospho-EGFR (Tyr1092) antibody.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Phospho-EGFR (Tyr1092) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

注意事项 (Note)

For research use only .