

产品名称: IKK beta Rabbit Monoclonal Antibody
产品货号: AMRe02148



产品概述 (Summary)

产品名称 (Production Name)	IKK beta Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	IKBKB IKBKB; IKKB; Inhibitor of nuclear factor kappa-B kinase subunit beta; I-kappa-B-kinase beta; IKK-B; IKK-beta; Ikbkb; I-kappa-B kinase 2; IKK2; Nuclear factor NF-kappa-B inhibitor kinase beta; NFKBIB
别名 (Alternative Names)	
基因 ID (Gene ID)	3551
蛋白 ID (SwissProt ID)	O14920.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000,IP 1:20-1:50
蛋白分子量 (Molecular Weight)	Calculated MW: 87 kDa; Observed MW: 87 kDa

产品名称: IKK beta Rabbit Monoclonal Antibody
产品货号: AMRe02148




研究背景 (Background)

The NF- κ B/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory I κ B proteins (1-3). Most agents that activate NF- κ B do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of I κ B (3-7). The key regulatory step in this pathway involves activation of a high molecular weight I κ B kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits.

研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Western blot analysis of IKK beta in HUVEC lysates using IKK beta antibody.

注意事项 (Note)

For research use only .