

产品名称: KRIT1 Rabbit Monoclonal Antibody
产品货号: AMRe02197



产品概述 (Summary)

产品名称 (Production Name)	KRIT1 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	KRIT1
别名 (Alternative Names)	CAM; CCM1
基因 ID (Gene ID)	889
蛋白 ID (SwissProt ID)	O00522.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000
蛋白分子量 (Molecular Weight)	Calculated MW: 84 kDa; Observed MW: 84 kDa

研究背景 (Background)

Component of the CCM signaling pathway which is a crucial regulator of heart and vessel formation and integrity .

产品名称: KRIT1 Rabbit Monoclonal Antibody
产品货号: AMRe02197




Negative regulator of angiogenesis. Inhibits endothelial proliferation, apoptosis, migration, lumen formation and sprouting angiogenesis in primary endothelial cells. Promotes AKT phosphorylation in a NOTCH-dependent and independent manner, and inhibits ERK1/2 phosphorylation indirectly through activation of the DELTA-NOTCH cascade. Acts in concert with CDH5 to establish and maintain correct endothelial cell polarity and vascular lumen and these effects are mediated by recruitment and activation of the Par polarity complex and RAP1B. Required for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction, and cell junction stabilization. Plays a role in integrin signaling via its interaction with ITGB1BP1; this prevents the interaction between ITGB1 and ITGB1BP1. Microtubule-associated protein that binds to phosphatidylinositol 4,5-bisphosphate (PIP2)-containing membranes in a GTP-bound RAP1-dependent manner. Plays an important role in the maintenance of the intracellular reactive oxygen species (ROS) homeostasis to prevent oxidative cellular damage. Regulates the homeostasis of intracellular ROS through an antioxidant pathway involving FOXO1 and SOD2. Facilitates the down-regulation of cyclin-D1 (CCND1) levels required for cell transition from proliferative growth to quiescence by preventing the accumulation of intracellular ROS through the modulation of FOXO1 and SOD2 levels.

研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Western blot analysis of KRIT1 in C6, 3T3, HeLa, K562 lysates using KRIT1 antibody.

注意事项 (Note)

For research use only .