

产品名称: RRP4 Rabbit Monoclonal Antibody
产品货号: AMRe02557



产品概述 (Summary)

产品名称 (Production Name)	RRP4 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	EXOSC2
别名 (Alternative Names)	p7; RRP4; SHRF; Rrp4p; hRrp4p
基因 ID (Gene ID)	23404
蛋白 ID (SwissProt ID)	Q13868.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000
蛋白分子量 (Molecular Weight)	Calculated MW: 33 kDa; Observed MW: 33 kDa

研究背景 (Background)

Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a

产品名称: RRP4 Rabbit Monoclonal Antibody
产品货号: AMRe02557




multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes. EXOSC2 as peripheral part of the Exo-9 complex stabilizes the hexameric ring of RNase PH-domain subunits through contacts with EXOSC4 and EXOSC7.


研究领域 (Research Area)

Epigenetics and Nuclear Signaling

图片 (Image Data)



Western blot analysis of RRP4 in rat spleen lysates using RRP4 antibody.



Western blot analysis of RRP4 in 3T3, HeLa lysates using RRP4 antibody.

产品名称: RRP4 Rabbit Monoclonal Antibody
产品货号: AMRe02557



注意事项 (Note)

For research use only .